

Anti-microbial and Anti-biofilm activity of leaf fruit and stem *Capparis spinose* extract against *Acinetobacter baumannii*.

Saeide Saeidi*, Elham Javadian, Fereshteh Javadian

Faculty of Sciences, Department of Biology, Science and Research Branch, Islamic Azad University, Kerman, Iran

Abstract

Background: Now-a-days, the increase in drug resistance among *Acinetobacter* is a major concern all over the world. The aim of this study was to investigate the antimicrobial and anti-biofilm activity of ethanolic extract and ethyl acetate of Leaf, Fruit and Stem *Capparis spinose* extract against *Acinetobacter baumannii*.

Methods: Twelve strains of *A. baumannii* were isolated from patients who referred to hospitals in Zabol. Antibiotic resistance pattern was determined by Kirby-Bauer method. Minimum inhibitory concentration and minimum bactericide concentration were determined by microdilution method.

Results: The results of the antibiotic resistance pattern showed that 100% of the samples were susceptible to CZ (Cefazolin)-GM (Gentamycin) - AZM (Azithromycin) antibiotics and only 16.6% of the samples were resistant to AM (Ampicillin) antibiotics. The results of this study showed that the lowest inhibitory concentration of the *C. spinose* ethanolic extract against *A. baumannii* was 25 ppm in which two bacterial strains were inhibited while the highest inhibitory concentration was 50 ppm and the highest concentration of fecundity was equal to 100 ppm.

Conclusion: Results of this study show the antimicrobial effects of good *C. spinose* biofilm on *A. baumannii*, which can be used as an effective treatment.

Keywords: *Acinetobacter baumannii*, Antibiotic resistance, Plant extract, *Capparis spinosa*.

Accepted on April 23, 2019

Introduction

Acinetobacterium is a gram negative bacterium and negative lactose that is increasingly recognized as an important factor in hospital infections such as the bacteremia and pneumonia associated with ventilator, surgical infections, secondary meningitis and urinary tract infections, especially in patients admitted to Special care departments. Its colony size is between 1 mm to 2 mm, no pigment and smooth to mucoid [1]. Blood agar usually causes hemolysis and grows at 44 °C. In the growth stage, the logarithmic form has a basil shape, but in the stagnant phase it is seen as coco-basil. It is a non-fermented bacterial group [2].

Various species of this bacterium are widely distributed in nature and can be separated from water, soil, human skin, food and sewage [3]. The genus of *Acintobacteria* consists of 23 species with a valid name and 10 anonymous species [4].

Acinetobacter baumannii has the ability to bind to all biological and non-biological levels, and form biofilms. This ability is an important pathologic property in many bacteria and promotes the development of antibiotic resistance and invasion of host immune system [5]. Being a community of microbial cells that are firmly attached to the surface, they are called biofilms with a microbe-derived polysaccharide matrix. Basically, bacteria create biofilms that resist antimicrobial

resistance to the host immune system and maintain the physical and chemical conditions suitable for growth, which makes biofilms resistant to adverse conditions. In the meanwhile, the synergistic and interphase relationships of biofilm bacteria are effective in their resistance to adverse conditions [6].

Populations of a species or different species of bacteria are adhered to each other or to living and non-living surfaces in the field of biofilm extracellular polysaccharides. Biofilm development is a complex process that requires bacterial communicative behaviours and, in contrast to single life, has many benefits for bacteria [7]. The formation of biofilms involves the reversible and irreversible initial binding to the surfaces, the formation of microcells, and, ultimately, the microcrystalline maturity associated with the formation of exopolysaccharides [8]. The spread of biofilm is the main cause of hospital infections because the bacteria that are attached to the surface and protected by external polysaccharide against various types of antibiotics and other environmental stresses, and biofilms increase the bacterial dispersion and bacterial pathology. Since these bacteria are resistant to various antibiotics, they now use plant extracts to eliminate them [9].

Capparis spinosa (locally called 'Al-Kabara') is one such plant stable to have highly diverse economic and pharmaceutical

value in various system of pharmaceutical like in Unani, Chinese, Ayurvedic and Greco-Arabi System of pharmaceutical. *C. spinosa* is well recognized with its usual name 'Capers' in various countries. In Iran, *C. spinosa* fruits and roots are generally apply to treat hemorrhoids [10,11].

Although the first snake grass habitat is not well known, today the range of herbs has expanded in a variety of Mediterranean and tropical climates to the Himalayan heights. In Iran, snake grass is found in most of the southern half of the country. Some species of the genus *Capparis* have been used in traditional medicine. It is used as a seasoning for peanuts, and its fruit is widely used in different cultures, which is one of the most important uses for pickles. Of course, the products of this plant may become susceptible in some people [12].

Other parts of this plant are used in the manufacture of medicines and cosmetics. Over the past four decades, snake grass has been considered for its economic importance as a crop in some European countries such as Greece and Italy [13]. Also, fruit and grass are used to treat diabetes, fungal infections, digestive tract infections and skin diseases caused by parasites, either orally or septa. Sometimes this plant is used to treat dryness and inflammation of the skin and to increase the blood flow to the skin as an ointment [14].

Materials and Methods

Separation of *A. baumannii*

Different strains of *A. baumannii* used in this research were isolated of urine specimens from hospitalized patients in Amiralmomenin Hospital of Zabol (Zabol-Iran). The 50 samples were cultured on agar Blood agar, BHI and agar Nitrite. To detect different species of Acinetobacter, the biochemical tests of urea-azcatalase and oxidase were performed.

Antibiotic susceptibility test was performed using a disk diffusion method in accordance with laboratory and clinical standards on the Mueller-Hinton agar medium. Streaking culture of bacterial suspensions of 0.5 McFarland were applied to the Muller Hinton Agar medium. In addition, the antibiotic discs were used including CZ (Cefazolin)-GM (Gentamycin)-AZM (Azithromycin)- AM (Ampicillin) (Mast, UK) [15].

Plant materials

The leaves, fruit, stem of *C. spinosa* were collected from the sistan region of Iran and they were dried at 25°C. 10 g grinded powders was soaked in 100 mL ethanol 95% and ethyl acetate, separately for one day (shaking occasionally with a shaker). After one day of dissolving process, materials were filtered. Then the filtrates were evaporated using rotary evaporator [16]. The broth microdilution method was used to determine MIC and MBC. Briefly, serial doubling dilutions of the extract were prepared in a 96-well microtiter plate ranged from 6.25 ppm to 100ppm.

Biofilm formation assay in presence of the biocides

At first, the OD (Optical Density) was measured (600 nm) by using an automated ELISA counter, then, the content of each well of the microplate was aspirated and each well was washed three times with 250 µL of sterile physiological saline. The remaining attached bacteria were fixed with 200 µL of 99% methanol per well and after 15 min all of the wells were emptied and left to dry. Then, each well was stained for 5 minute with 0.2 mL of 2% crystal violet. Excess stain was rinsed off by washing the plate slowly with distilled water. After the plate was air dried, the dye bound to the adherent cells was resolubilized with 160 mL of 33% (v/v) glacial acetic acid per well. The OD of each well was measured at 492 nm by using an automated ELISA counter.

Statistical analyses

The growth was compared at each experiment using analysis of variance (ANOVA) repeated measures (SPSS-25.0 for Windows). The level of statistical significance was set at $P < 0.01$.

Results

The results of the antibiotic resistance pattern showed that 100% of the samples were susceptible to CZ (Cefazolin)-GM (Gentamycin)-AZM (Azithromycin) antibiotics and only 20% of the samples were resistant to AM (Ampicillin) antibiotics (Table 1). The results of this study showed that the lowest inhibitory concentration of the *C. spinosa* ethanol extract against *A. baumannii* was 25 ppm, in which two bacterial strains were inhibited, while the highest inhibitory concentration was 50 ppm and the highest concentration of fecundity is equal to 100 ppm (Table 2). The results of the study of *C. spinosa* ethanol extract showed that the lowest inhibitory concentration was 25 ppm, while the highest inhibitory concentration was 100 ppm, which was inhibited by a strain of bacteria.

Table 1. Patten antibiotic of *A. baumannii*.

Antibiotic	Resistant (%)	Intermediate (%)	Sensitive (%)
AMC	1 (10%)	1 (10%)	8 (80%)
AM	2 (20%)	3 (30%)	5 (50%)
GM	0 (0%)	0 (0%)	0 (100%)
CZ	0 (0%)	0 (0%)	0 (100%)
AZM	0 (0%)	0 (0%)	0 (100%)

The results of this study showed that the lowest inhibitory concentration of ethanolic extract of grass snake was 12.5 ppm, two strains were inhibited in this concentration, while the highest inhibitory concentration of ethanolic extract was 50 ppm, 5 of which in this concentration of inhibition.

The lowest inhibitory concentration of the gum extract ethyl acetate was 6.25 ppm. Two strains were inhibited at this

Anti-microbial and Anti-biofilm activity of leaf fruit and stem Capparis spinose extract against Acinetobacter baumannii

concentration, while the highest germination concentration of the ethyl acetate extract was 100 ppm; one strain was inhibited in this concentration (Table 3).

Table 2. Minimum inhibitory concentration and minimum bactericide concentration of *C. spinose* leaves against *Acinetobacter* (ppm).

Bacterium sample	Ethanol extract leaf		Ethyl acetate extract leaf	
	MIC	MBC	MIC	MBC
1	50	25	50	25
2	50	25	50	25
3	100	50	100	50
4	100	50	50	25
5	100	50	50	25
6	100	50	50	25
7	100	50	50	25
8	100	50	50	25
9	100	50	50	25
10	100	50	50	25
11	100	50	50	25
12	100	50	100	100

Table 3. Minimum inhibitory concentration and minimum bactericide concentration of *C. spinose* fruit against *Acinetobacter* (ppm).

Bacterium samples	Ethanol extract fruit		Ethyl acetate extract fruit	
	MIC	MBC	MIC	MBC
1	25	12.5	50	25
2	50	25	25	12.5
3	25	12.5	50	25
4	100	50	50	25
5	50	25	12.5	6.25
6	100	50	50	25
7	50	25	50	25

Table 5. Results of biofilm OD at different concentrations of ethanol extract of leaf (ppm).

Variables	Concentration (ppm)							Bacterium sample
	3.1	6.25	12.5	25	50	100		
Control biofilm any extract								
1.256	0.93	1.021	1.553	0.165	0.236	0.221	1	
1.562	1.266	1.24	1.513	0.727	0.201	0.109	2	
1.195	1.087	1.185	0.98	1.38	0.202	0.248	3	
1.428	1.307	1.252	1.087	0.343	0.299	0.158	4	
1.346	1.197	1.192	0.902	0.55	0.167	0.134	5	

8	100	50	25	12.5
9	100	50	50	25
10	50	25	12.5	6.25
11	100	50	50	25
12	50	25	100	50

The results showed that the minimum inhibitory concentration of ethanolic extract of root extract was 50 ppm; all 12 strains were inhibited in this concentration, while the lowest inhibitory concentration of root extract acetate extract was 25 ppm, of which 7 strains were inhibited in this concentration (Table 4).

Table 4. Minimum inhibitory concentration and minimum bactericide concentration of *C. spinose* stem against *Acinetobacter* (ppm).

Bacterium sample	Ethyl acetate extract stem		Ethanol extract stem	
	MIC	MBC	MIC	MBC
1	50	25	50	50
2	50	25	50	50
3	100	50	100	50
4	100	50	100	50
5	50	25	100	50
6	50	25	100	50
7	50	50	100	50
8	50	25	100	50
9	50	25	100	50
10	50	50	100	50
11	50	25	100	50
12	100	100	100	50

The results of the OD results for biofilm formation showed that at higher concentrations, the inhibitory effect of biofilm formation was lower (Tables 5 and 6).

1.452	1.11	1.263	1.213	0.462	0.238	0.194	6
1.325	1.199	1.264	0.897	0.345	0.198	0.165	7
1.121	0.957	1.462	0.993	0.354	0.18	0.203	8
1.248	0.98	1.144	0.571	0.436	0.186	0.157	9
1.359	1.166	0.902	1.124	0.36	0.246	0.168	10
1.526	1.19	0.907	0.732	0.278	0.204	0.153	11
1.369	1.127	1.153	0.566	0.697	0.637	0.148	12

Table 6. Results of biofilm OD at different concentrations of ethyl acetate extracts of leaf (ppm).

Control biofilm any extract	Concentration (ppm)						Bacterium sample
	3.1	6.25	12.5	25	50	100	
1.489	1.338	1.309	1.196	0.262	0.21	0.202	1
1.925	1.708	1.585	1.313	0.5	0.268	0.169	2
1.759	1.55	1.479	1.143	0.407	0.299	0.181	3
1.895	1.581	1.573	0.901	0.397	0.283	0.177	4
1.855	1.664	1.552	1.47	0.513	0.28	0.274	5
1.639	1.568	1.5	1.449	0.729	0.249	0.359	6
1.728	1.68	1.3	1.088	0.394	0.369	0.196	7
1.658	1.529	1.339	1.276	0.186	0.393	0.208	8
1.432	1.137	1.36	1.54	0.45	0.282	0.225	9
1.563	1.335	1.475	1.336	0.434	0.276	0.175	10
1.697	1.379	1.446	0.995	0.295	0.242	0.204	11
1.658	1.317	0.768	0.935	0.215	0.263	0.192	12

Discussion

The results of the antibiotic resistance pattern showed that 100% of the samples were susceptible to CZ-GM (Gentamycin)-AZM antibiotics and only 16.6% of the samples were resistant to AM (Ampicillin) antibiotics.

In the study of Goodarzi et al. [17] on antibiotic resistance pattern of *A. baumannii* in Tehran, the results showed that the resistance strains to antibiotics were: 103 (95.4%) to ceftazidime, 108 (100%), 100 (6/92%) to ciprofloxacin, 103 (95.4%) to piperacillin-tazobactam, 44 (40.7%) to gentamicin, 87 (80.6%) to tetracycline and 1 (0.9%) to clostin.

The study of Jafari et al. [18] examined the antibiotic resistance pattern of *A. baumannii* in Fars province and the results showed that the percentages of *Acinetobacter* isolates of amphenem, moropenem and ciprofloxacin in serial dilution method were 40.9%, 60%, and 77.7%; respectively. In a study by Simhon et al. [19] sensitivity to ampicemin decreased from 98.1% in 1990 to 64.1% in 2000, and the sensitivity to ciprofloxacin decreased from 50.5% to 13.1%. A study in UK was conducted by Caroline and colleagues in 2002, and results

showed that 46% of isolates were resistant to ciprofloxacin and 2% of isolates were resistant to amipenem [20].

In a study conducted in Tehran in 2009 by Boromand et al., 53.4% of the samples were resistant to ciprofloxacin and 24.6% of them were resistant to amipenem [21]. Antibiotic resistance patterns among hospital pathogens may vary widely from one country to another or in different regions of a country. Previous reports from around the world suggest that 30-83.9% of Acetone is resistant to multiple medicines [22]. On the other hand, Ferreira et al. found that 68% were resistant to carbapenems [23].

Peymani et al. [24] found that in 81% of isolates in antibiotic classes beta-lactam, aminoglycosides and quinolones, there were complete or intermediate drug resistance in 2012, and their percentage of resistance to ampicemin and maropenem were 68% and 69%, respectively. The results of the OD results for biofilm formation showed that at higher concentrations, the inhibitory effect of biofilm formation was lower.

In 1394, the effect of alcoholic essence of *Satureja khuzestanica* on the expression of the gene associated with Biofilm *A. baumannii* was investigated. Due to the inhibitory

effect of the Khuzestan Sourifolia essential oil on the Biofilm *A. baumannii* gene, it would probably be possible to use this essential oil as a supplement [25]. The results of this study showed that the lowest inhibitory concentration of the *C. spinosa* ethanol extract against *A. baumannii* was 25 ppm, in which two bacterial strains were inhibited, while the highest inhibitory concentration was 50 ppm and the highest concentration of fecundity is equal to 100 ppm. The results of the study of *C. spinosa* ethanol extract showed that the lowest inhibitory concentration was 25 ppm, while the highest inhibitory concentration was 100 ppm, which was inhibited by a strain of bacteria.

The study of Abd Razik [26], antibacterial effects of different concentrations ranging from (125-1000 mg/ml) of hexane and methanol juice of *C. spinosa* flowers was determined by using agar well diffusion method in clinical strains. Methanol juice of *C. spinosa* flowers was the most active than hexane juice. The activity of *C. spinosa* determined by measuring inhibition region as following: *Lactobacillus sp.*, *Escherichia coli*, *Streptococcus sp.* and *Staphylococcus aureus* demonstrate inhibition region at concentration of 125 -1000 mg/ml. *Klebsiella sp.*, *Pseudomonas aeruginosa* and *Enterococcus sp.* showed inhibition region of 500- 1000 mg/ ml. it was no inhibition region for *Proteus sp.* Hexane juice of *C. spinosa* flowers was less active than methanol juice against examine bacteria. *Lactobacillus sp.*, *S. aureus* and *Streptococcus sp.* demonstrate inhibition region at concentrations 150- 1000 mg/ml while *Escherichia coli*, *S. aureus* and *Klebsiella sp.* demonstrate sensitivity at concentration 1000 mg/ml. There was no inhibition region for *P. aeruginosa*, *Enterococcus sp.* [26]. Ibrahim [27] investigated the antimicrobial acting of the flower extract of *C. spinosa*, gel and leaf extract of Aloe vera were examine against isolated bacterial of skin infection *S. aureus*, *S. epidermidis*, *S. pyogenes*, *P. aeruginosa* and *E. coli*. Antimicrobial susceptibility examine demonstrated that the *C. spinosa* was 100% effective versus gram positive isolates and 90% acting against gram negative isolates, A. vera leaf was 20% effective versus the entire tested gram positive as well as 15% effective versus gram-negative isolates. *Capparis spinosa* showed prominent antibacterial activity versus gram negative and positive.

Benakashani et al. [28] studied the antimicrobial effect of nanoparticles produced *C. spinosa* by disc diffusion method against different pathogenic bacteria such as *E. coli*, *Salmonella typhimurium*, *S. aureus* and *Bacillus cereus*. The results showed that synthesized silver nanoparticles have an excellent antimicrobial property and a high antimicrobial acting compared to the ionic silver.

Biofilm is a structural society and a social expression of microorganisms, which are placed in a space between them by the surface of the enamelled polyester produced by them and connected to one surface. Biofilm formation is an active process involving the initial binding, the formation of microcolony, the production of extracellular polymeric material, followed by growth and maturation, and ultimately

thinning and releasing the cell from biofilms [29]. Since their antibiotic resistance is increasing, it requires new treatments.

Bahador et al. [30] studied effect of Satureja khuzestanica essence, on the expression level of *bap* gene in *A. baumannii*. The results showed by reducing *bap* gene expression. They hope in future be used it to the clinic with a wider range as a complementary therapy and also for surgery operation.

José Alves et al. [31] reported the *Acinetobacter baumannii* was the microorganism with the lowest susceptibility to mushroom extracts inhibitory effect on biofilm production (highest inhibition—almost 29%, by *Russula delica* extract). David Alejandro et al. [32] investigated the effect of sub-lethal concentrations of MBC of the extract, by inhibiting the formation of biofilms. The result show that concentrations of 25%, 50% and 75% of the MBC (1.87, 3.74 and 5.61 mg/mL, respectively), caused biofilms inhibitions of 19.5%, 40.8% and 36.0%, respectively.

Conclusion

The results of this study show the anti-microbial effects of good *C. spinosa* biofilm on *A. baumannii*, which can be used as an effective treatment. The results of this study show the good antimicrobial effects of the extract *C. spinosa* on the pathogenic bacteria. At first, we resolved total phenolic and total flavonoid extent of each extract of fruits and roots.

Conflict of Interests

The authors declare no conflict of interest.

Funding/Support

This study was supported by the Institute of Plant Biotechnology, University of Zabol, Zabol, Iran.

References

1. Qi C, Scheetz MH, Malczynski M. Characterization of *Acinetobacter baumannii* genotypes recovered from patients with repeated colonization or infection. *Diagn Microbiol Infect Dis.* 2009; 65: 1-6.
2. Khoshriahi N, Sharifi M. Isolation of carbapenem resistant *Acinetobacter baumannii* (CRAB) strains from patients and equipments of Intensive care units (ICUs) at Qazvin between 2005-2006. *Iran J Med Microbiol.* 2007;1: 33-84.
3. Glew RH, Moellering RC, Kunz LJ. Infections with *Acinetobacter alcoaceticus* (*Herellea vaginicola*): Clinical and laboratory studies. *Medicine.* 1977; 56: 79-97.
4. Visca P, Seifert H, Towner KJ. *Acinetobacter* infection: An emerging threat to human health. *IUBMB Life.* 2011; 63: 1048-54.
5. Zarrilli R, Crispino M, Bagattini M, et al. Molecular epidemiology of sequential outbreaks of *Acinetobacter baumannii* in an intensive care unit shows the emergence

- of carbapenem resistance. *J Clin Microbiol.* 2004; 42: 946-53.
6. Lear G, Lewis GD. *Microbial biofilms: Current research and applications*: Horizon Scientific Press. 2012.
 7. Gubner R, Beech I. The effect of extracellular polymeric substances on the attachment of *Pseudomonas NCIMB 2021* to AISI 304 and 316 stainless steel. *Biofouling.* 2000; 15: 25-36.
 8. Der Mei HC, Busscher HJ. Physico-chemistry of initial microbial adhesive interactions—Its mechanisms and methods for study. *FEMS.* 1999; 23: 179-230.
 9. Bos R, Van, Kojic EM, Darouiche RO. *Candida* infections of medical devices. *CMR* 2004; 2: 255- 67.
 10. Azaizeh H, Fulder S, Khalil K, et al. Ethnomedicinal knowledge of local Arab practitioners in the Middle East Region. *Fitoterapia.* 2003; 74: 98-108.
 11. Miraldi E, Ferri S, Mostaghimi V. Botanical drugs and preparations in the traditional medicine of West Azerbaijan (Iran). *J Ethnopharmacol.* 2001; 75:77-87.
 12. Bahador A, Saghii H, Ataee R, et al. The study of inhibition effects *Satureja khuzestanica* essence against gene expression *baumannii* with Real time PCR Technique. *Iran J Med Microbiol.* 2015; 9: 42-49.
 13. Costerton JW, Lewandowski Z, Caldwell DE, et al. *Microbiol Biofilm.* *Ann Rev Microbiol.* 1995; 49: 711-45.
 14. Maria José Alves, Isabel C. F. R. Ferreira, Inês Lourenço, et al. Wild mushroom extracts as inhibitors of bacterial biofilm formation pathogens. 2014; 3: 667-679.
 15. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing (2013), 16th International supplement. CLSI document M100-S21.
 16. Hanafy MS, Hatem ME. Studies on the antimicrobial activity of *Nigella sativa* seed (black cumin). *J Ethnopharmacol.* 1991; 34: 275-278.
 17. Goodarzi H, Basharati M, Zarrin Kalam Moghadam M. Investigation of drug resistance of *Acinetobacter bomani* isolated from patients admitted to Imam Reza Hospital of Tabriz in 2013. *Research in Medical Sciences.* 2014; 38: 106-110.
 18. Jafari S, Najafipour S, Kargar M, et al. Phenotypic study of drug resistance *Acinetobacter Bohemian* with multiple drug resistance. *J of Fasa University of Med Sci.* 2012; 2.
 19. Simhon A, Rahav G, Shazberg G, et al. *Acinetobacter baumannii* at a Tertiary-Care Teaching Hospital in Jerusalem. *J Clin Microbiol.* 2001; 39: 389-91.
 20. Henwood C, Gatward T, Warner M, et al. Antibiotic resistance among clinical isolates of *Acinetobacter* in the UK, and in vitro evaluation of Tigecycline) GAR-936 (: *Antimicrob chemother.* 2004; 49: 479-487.
 21. Boromand M, Akyani M, Sheikhvatan R, et al. Evaluation of antimicrobial resistance of *Acinetobacter baumannii* to Im-ipenem, Ciprofloxacin and Ceftazidim using E test. *Iranian J Publ Health.* 2009; 2: 130-133.
 22. Kwmpf M, Rolain JM. Emergence of resistance to carbapenems in *Acinetobacter baumannii* in Europe: clinical impact and therapeutic options. *Int J Antimicrob Agents.* 2012; 39:105-114.
 23. Ferreira AE, Marchetti DP, Cunha GR, et al. Molecular characterization of clinical multiresistant isolates of *Acinetobacter sp.* From hospitals in Porto Alegre, State of Rio Grande do Sul, Brazil. *Rev Soc Bras Med Trop.* 2011; 44: 725-30.
 24. Peymani A, Farajnia S, Nahaei MR, et al. Prevalence of class 1 integron among multidrugresistant *Acinetobacter baumannii* in Tabriz, North West of Iran. *Pol J Microbiol.* 2012; 61: 57-60.
 25. Bahador A, Saghii H, Ataee R, et al. The Study of Inhibition Effects *Satureja khuzestanica* Essence against Gene Expression *baumannii* with Real time PCR Technique. *Iranian J Med Microbiol.* 2015; 9: 42-9.
 26. Abd Razik B. The Study of Antibacterial Activity of *Capparis spinosa* Flowers *Al-Mustansiriyyah J. Sci.* 2011; 5.
 27. Ibrahim MS. Evaluation of anti-bacterial activity of *Capparis spinosa* (Al-Kabara) and *Aloe vera* extracts against Isolates Bacterial Skin Wound Infections in -vitro and in-vivo. *Kufa Journal For Veterinary Medical Sciences.* 2012; 3.
 28. Benakashani F, Allafchian AR, Jalali SAH. Biosynthesis of silver nanoparticles using *Capparis spinosa* L. leaf extract and their antibacterial activity. *Karbala Int J Mod Sci* 2014; 2: 251-258.
 29. Hernández-Marín David Alejandro, Muñoz-Ortega Martín Humberto, et al. Inhibition of *Acinetobacter baumannii* biofilm formation by methanolic extract of *nothoscordum bivalve*. *Scientific Research* 2018; 8.
 30. Alcantara M, Morales M, Carnes J. Food allergy to caper (*Capparis spinosa*). *J Investig Allergol Clin Immunol.* 2013; 23: 67-69.
 31. Yang T, Wang C, Liu H, et al. A new antioxidant compound from *Capparis spinosa*. *Pharmaceutical Biol.* 2010; 48: 589-94.
 32. Zhou H, Jian R, Kang J, et al. Anti-inflammatory effects of caper (*Capparis spinosa* L.) fruit aqueous extract and the isolation of main phytochemicals. *J Agric Food Chem.* 2010; 58: 12717-21.

*Correspondence to

Saeide Saeidi

Faculty of Sciences

Department of Biology

Islamic Azad University

Kerman

Iran