

Diagnostic and prognostic role of Procalcitonin in sepsis in a tertiary care hospital.

Sunil Kumar Nanda^{1*}, Asha Dinakaran², Sandhya Bhat K³, Ravichandran⁴, Reba Kanungo⁵

¹Department of Biochemistry, Pondicherry Institute of Medical Sciences, Affiliated to Pondicherry University, Pondicherry.

²Department of Biochemistry, Pondicherry Institute of Medical Sciences, Affiliated to Pondicherry University, Pondicherry.

³Department of Clinical Microbiology, Pondicherry Institute of Medical Sciences, Affiliated to Pondicherry University, Pondicherry.

⁴Department of Biostatistics, Pondicherry Institute of Medical Sciences, Affiliated to Pondicherry University, Pondicherry.

⁵Department of Clinical Microbiology, Pondicherry Institute of Medical Sciences, Affiliated to Pondicherry University, Pondicherry.

Abstract

Background: Bacterial sepsis is one of the common causes of morbidity and mortality in patients admitted to any tertiary care hospital. Early diagnosis of bacterial sepsis is a challenge to the clinician and the laboratory. Blood culture is the gold standard for the diagnosis of bacterial sepsis. Isolating the bacteria and determining its antibiotic susceptibility takes a minimum of 48 hours. There is a constant search for biochemical markers which might rapidly indicate bacterial sepsis.

Aim: The aim of the study was to evaluate the diagnostic and prognostic role of procalcitonin in bacterial sepsis.

Subject and methods: Procalcitonin (PCT) levels were determined in 136 patients admitted to the hospital with symptoms of sepsis. PCT was measured in serum samples obtained from the patients at the same time when blood was drawn for culture.

Results: Out of the 136 patients clinically diagnosed with sepsis, 14 patients had a positive blood culture (10.3%). Procalcitonin levels of >0.5 ng/ml was considered as positive. Out of the 14 patients with a positive blood culture, 12 had PCT levels greater than 0.5 ng/ml. ROC analysis revealed maximum sensitivity of 64% and a specificity of 65% with a PCT value of 3.25 ng/ml. The mean PCT values in survivors were lower when compared to PCT values in non survivors.

Conclusion: Markers for rapid diagnosis of sepsis is required to institute rapid and appropriate management. Procalcitonin level is one such promising target marker for early diagnosis of sepsis.

Keywords: Procalcitonin, Bacterial sepsis, Blood culture.

Accepted November 26, 2015

Introduction

Bacterial sepsis is associated with high morbidity and mortality. Sepsis is the second most common cause of death after myocardial infarction in patients admitted to intensive care units. Mortality due to sepsis is as high as 25-35% and is higher in patients with septic shock [1]. Early diagnosis is critical in management of patients with bacterial sepsis since appropriate antibiotics can be initiated immediately. Blood culture remains the gold standard for diagnosis of bacterial sepsis. However a major limitation is unavailability of results within 48 hrs. Prior antibiotic therapy often results in a negative blood culture there by negating a confirmatory diagnosis of sepsis. Skin colonizers often confound a culture outcome in patients presenting with fever due to non-bacterial causes, resulting in unnecessary antibiotic therapy. Several clinical markers

of infection have been recommended but most are found to be non-specific since they can be positive in systemic inflammation of non-infectious origin. Hematological markers of infection like total and differential counts may also be non-specific [2,3,4].

There is a continuous search for biomarkers of sepsis. Some of the biomarkers that have been evaluated include lactate, interleukins, C reactive protein (CRP) and procalcitonin [4,5,6]. C reactive protein is most widely used acute phase reactant and a sensitive marker of inflammation. It cannot differentiate bacterial sepsis from other causes of inflammation. CRP gets elevated only 24 to 48 h after the infection is initiated, hence cannot be a rapid indicator [7,8]. Reports of Procalcitonin (PCT) include its role in diagnosis of bacterial sepsis, determining the severity of sepsis and in determining the duration of antibiotic

administration in children and adults [2,9]. This study was designed to determine procalcitonin levels in patients with sepsis and its correlation with blood culture outcome.

Materials and Methods

The study was a retrospective study. 136 adult patients (>18 years of age) admitted to a tertiary care hospital between November 2012 to May 2014 were included in the study. The sample size was calculated by taking into account the prevalence of suspected cases of sepsis, sensitivity and specificity of Procalcitonin (PCT). The study was approved by the institutional ethical committee.

Materials

Data on clinical features, laboratory investigations, blood culture and PCT values were retrieved from the medical records department of the institute. A diagnosis of sepsis was made based on the recommendations of the American College of Chest Physicians (ACCP) which included presence of any of the following -2 or more of the features along with suspected or proven source of infection: Temp >38°C (100.4°F) or <36°C (96.8°F), Heart Rate >90, Respiratory Rate >20 or PaCO₂ <32 mmHg, WBC >12,000/mm³, <4,000/mm³, or >10% bands. Patients with cardiogenic shock, small cell lung carcinoma, medullary carcinoma of thyroid, major trauma, major surgical intervention, severe burns were excluded from the study, as PCT is nonspecifically elevated in these conditions.

Methods

Blood culture of all 136 patients was done by automated BacT/Alert system. Procalcitonin was measured in Roche e411 Electrochemiluminescence (ECLIA) automated analyzer using PCT kit from B.R.A.H.M.S Diagnostica, Berlin, Germany. According to the manufacturers, a value of PCT >0.5 ng/ml was taken as pathological, 0.5 to 2 ng/ml indicated that systemic infection could not be ruled out, 2 to 10 ng/ml indicated greater chances of sepsis and a value of PCT above 10 ng/ml indicated severe bacterial sepsis.


Statistical Analysis

Validation of PCT was done by calculating the sensitivity,

specificity, positive predictive value, negative predictive value. The cut off value of PCT which gave maximum sensitivity and specificity were calculated by Receiver operator curve (ROC) analysis.

Results

Blood culture positivity and the bacteria isolated in different ranges of PCT values are given in Table 1. The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) of PCT by taking a cut off of 0.5 ng/ml or more as positive for sepsis are given in Table 2. Receiver operator curve (ROC) analysis was done to determine the cut off value of PCT for which maximum sensitivity and specificity was obtained. ROC analysis is illustrated in Figure 1 which indicates a maximum sensitivity of 64% and specificity of 65% at PCT value of 3.25 ng/ml.



Diagonal segments are produced by ties.

Figure 1: Receiver operator curve analysis of PCT levels in patients with sepsis with sensitivity on Y axis and specificity on X axis.

Table 1: Serum PCT values, blood culture results in 136 patients of sepsis.

PCT Values	<0.5 ng/ml	0.5 to 1.9 ng/ml	2 to 10 ng/ml	>10 ng/ml
Total number of patients in number and percentage	35 (26%)	37(27%)	27(20%)	37 (27%)
Blood culture positives in patients in number and percentage	2 (14%)	2 (14%)	3 (22%)	7(50%)

Bacteria grown

E. coli, CONS

K. pneumonia,
A. baumannii

S. aureus,
Acinetobacter,
E. coli

M. morgani,
P. aeruginosa,
E. coli, *Acinetobacter*
Escherichia coli
Gram negative bacilli
P. aeruginosa

E. coli: *Escherichia coli*, *CONS*: *Coagulase negative Staphylococcus aureus*, *K. pneumonia*: *Klebsiellapneumoniae*, *A. baumannii*: *Acinetobacterbaumanni*, *S. aureus*: *Staphylococcus aureus*, *M. morgani*: *Morganellamorganii*, *P. aeruginosa*: *Pseudomonas aeruginosa*

Table 2: Sensitivity, specificity, PPV, NPV of PCT.

Sensitivity (True positives)	85.7%
Specificity (True negatives)	25.4%
PPV	11.7%
NPV	93.9%

Discussion

PCT is the precursor for the hormone calcitonin. Calcitonin has a metabolic role in calcium homeostasis. Though PCT is synthesized in the thyroid and pulmonary cells in normal conditions, all tissues throughout the body have the potential to express PCT [10]. Procalcitonin has been studied as a diagnostic marker in many conditions like fever of unknown origin, meningitis, respiratory tract infections, urinary tract infections, burns and blood stream infections. PCT has been studied as prognostic marker with high values indicating bacterial load and severity of sepsis. The concept of PCT clearance has been used as an indicator of recovery [11].

In the present study, by taking a cut off of >0.5 ng/ml as an indicator of sepsis, the sensitivity, specificity, PPV and NPV was found to be 85.7%, 25.4%, 11.7% and 93.9% (Table 2) and by taking a cut off of >2 ng/ml, the sensitivity, specificity, PPV and NPV was found to be 71.4%, 56.6%, 15.9% and 94.5%. Sinha M et al., observed a sensitivity of 90% and specificity of 84% for a cut off of 0.5 ng/ml where as a sensitivity of 85.7% and specificity of 94.7% was noted with cut off of 2 ng/ml. They concluded that PCT assay may avoid unwarranted antibiotic usage [12]. Harbarth et al., observed a sensitivity of 97% and specificity of 78% when a cut off of 1.1 ng/ml was used to diagnose systemic inflammatory response syndrome [SIRS] with a conclusion that PCT appeared to be a promising indicator of sepsis in newly admitted, critically ill patients and PCT values complemented with the clinical signs and routine laboratory parameters, suggestive of severe infection [5]. Sudhir U et al., observed that a sensitivity of 94% had a significant association between serum PCT and Sequential Organ Failure Assessment (SOFA) score [13].

Out of the 136 cases, 90 were males and 46 were females. The mean PCT values in males were 15.9 ng/ml and the mean PCT values in females were 13.9 ng/ml. The difference was not statistically significant. Out of the 136 cases, 14 patients had a positive blood culture. Commonest bacteria grown were *Escherichia coli* in 4 out of 14 culture positive cases (Table 1 and Figure 2). Sucilathangam G et al., observed *Acinetobacter* as the commonest organism grown in 5 out of 14 culture positive cases [14]. Endotoxins present in the bacterial cell wall induce the production of PCT from the parenchymal tissues. The defense response to infection also contributes to increase in PCT levels in bacterial infections. The parenchymal cells do not have the ability to convert PCT to calcitonin leading on to its increased levels in circulation [16,17]. Sinha M et al., observed gram positive cocci as the commonest isolates [12]. Findings of Wang H et al., showed *Escherichia coli*




Figure 2: Bacteria isolated indicating the most common bacteria isolated in culture was *E.coli* followed by *Acinetobacter*, *Pseudomonas aeruginosa*, *staphylococcus aureus*, *Kliebsiella pneumonia* and *morganella morganii*.

to be predominant isolate among gram negative organisms similar to the present study. Coagulase-Negative *Staphylococcus* and *Enterococcus* were the common gram positive isolates [18]. In the present study, we observed that out of 14 isolates, 12 were gram negative and only 2 isolates were gram positive. A possible correlation between gram negative endotoxins and PCT levels needs to be explored. The mean PCT values of patients with gram positive isolates and gram negative isolates were 2.9 ng/ml and 31 ng/ml but the difference in the values in the two groups were not statistically significant. (Table 3)

ROC analysis revealed, maximum sensitivity of 64% and maximum specificity of 65% for a PCT value of 3.25 ng/ml (Figure 1). In a study done by Riedel S et al., maximum sensitivity and specificity was observed for a much lower value of PCT at 0.1475 ng/ml [15].

The mean PCT value of those who succumbed to the infection (23 in number) was 30.6 ng/ml whereas the mean value in survivors was 12 ng/ml (Table 4). Our findings suggest that high PCT values can indicate high mortality rate when compared to patients with lower values, a finding similar to Jensen J.U et al., who observed that high PCT levels is an independent predictor of mortality [19]. On the other hand, Ruiz-Alvarez et al., observed that PCT did not predict mortality whereas CRP did [20]. Pettila et al., observed that there was statistically significant difference in the PCT values measured on first and second days in patients who survived than patients who did not survive [21].

Table 3: PCT values in gram positive and gram negative isolates.

Isolate	Mean PCT values (ng/ml) ± Standard error of mean	P value
Gram negative isolate	31 ± 10	0.306
Gram positive isolate	2.9 ± 2.5	

Table 4: PCT values in survivors and non survivors.

	Mean PCT values (ng/ml) ± Standard error of mean	P value
Survivors	12 ± 2.3	0.042*
Non Survivors	30.6 ± 8.3	

Table 5: PCT values in patients with less than 10 days duration of hospitalization and more than 10 days of hospitalization.

	Mean PCT values (ng/ml) ± Standard error of mean	P value
Patients with less than 10 days duration of hospitalization	15.7 ± 3.3	0.79
Patients with more than 10 days duration of hospitalization	14.5 ± 3.5	

It was observed that the mean PCT values of patients (80 in number) who were hospitalized for less than 10 days was 15.7 ng/ml whereas mean PCT values of patients hospitalized for more than 10 days was 14.5 ng/ml (Table 5). Two patients were culture positive but had PCT values within normal range. The isolate from blood of one of the patients was CONS, a normal skin commensal which might not have induced the production of PCT. In another case, a bacterium grown was *E. coli* and PCT was negative. The reason could have been initiation of empiric antibiotic prior to blood culture. It was observed that as the level of PCT increased, the chances of blood culture positivity increased.

Conclusion

Bacterial sepsis is often a medical emergency requiring intensive management with specific antibiotic therapy and other supportive measures. A delay in detection of sepsis may lead to poor outcome. Rapid sepsis markers in the blood will help in overcoming the limitations of confirmation by blood culture. Procalcitonin levels can play a major role in not only detecting sepsis but also to monitor progress or predict outcome.

References

1. Chaudhury A, Rao TV. Bacteraemia in a tertiary care urban hospital in south India. *Indian J Pathol Microbiol* 1999; 42: 317-320.
2. Castelli GP, Pognani C, Meisner M, Stuaní A, Bellomi D, Sgarbi L. Procalcitonin and C reactive protein during systemic inflammatory response syndrome, sepsis and organ dysfunction. *Crit Care* 2004; 8: 234-242.
3. Muller B, Schuetz P, Trampuz A. Circulating biomarkers as surrogates for bloodstream infections. *Int J Antimicrob Agents* 2007; 30: S16-23.
4. Muller B, Becker KL, Schachinger H, Rickenbacher PR, Huber PR, Zimmerli. Calcitonin precursors are reliable markers of sepsis in a medical intensive care unit. *Crit Care Med* 2000; 28: 977-983.
5. Harbarth S, Holekova K, Froidevaux C, Pittet D, Ricou B, Grau GE. Diagnostic value of procalcitonin, interleukin 6 and interleukin 8 in critically ill patients admitted with suspected sepsis. *Am J Respir Crit Care Med* 2001; 164: 396-402.
6. Nanda SK and Suresh DR. Plasma Lactate as prognostic marker of septic shock with acute respiratory distress syndrome. *Indian Journal of Clinical Biochemistry* 2009; 24: 433-435.

7. PCNg. Diagnostic markers of infection in neonates. *Arch Dis Child Fetal Neonatal* 2004; 89: F229-F235.
8. Weinschenk NP, Farina A, Bianchi DW. Premature infants respond to early-onset and late-onset sepsis with leukocyte activation. *J Pediatr* 2000; 137: 345-350.
9. Ali AM, Walid FE, Abdelaziz SS. Procalcitonin Versus C-Reactive Protein in Neonatal Sepsis. *J Immunol Infect Dis* 2014; 1: 1-5.
10. Kenneth LB, Snider R, Nylén ES. Procalcitonin in sepsis and systemic inflammation: a harmful biomarker and a therapeutic target. *British Journal of Pharmacology* 2010; 159: 253-264.
11. Mehanic S, Baljic S. The Importance of Serum Procalcitonin in Diagnosis and Treatment of Serious Bacterial Infections and Sepsis. *Mater Socio med* 2013; 25: 277-281.
12. Sinha M, Desai S, Mantri S, Kulkarni A. Procalcitonin as an adjuvant marker in sepsis. *Indian Journal of Anaesthesia* 2011; 55: 266-270.
13. Sudhir U, Venkatachalaiah RK, Kumar TA, Rao MY, Kempegowda P. Significance of serum procalcitonin in sepsis. *Indian J Crit Care Med* 2011; 15: 1-5.
14. Sucilathangam G, Amuthavalli K, Velvizhi G, Ashihabegum M.A, Jeyamurugan T, Palaniappan N. Early Diagnostic Markers for Neonatal Sepsis: Comparing Procalcitonin (PCT) and C-reactive protein (CRP). *Journal of Clinical and Diagnostic Research* 2012; 6: 627-631.
15. Riedel S, Melendez JH, An AT, Rosenbaum JE, Zenilman JM. Procalcitonin as a Marker for the Detection of Bacteremia and Sepsis in the Emergency Department. *Am J Clin Pathol* 2011; 135: 182-189.
16. Linscheid P, Seboek D, Schaer DJ, Zulewski H, Keller U. Expression and secretion of procalcitonin and calcitonin gene-related peptide by adherent monocytes and by macrophage-activated adipocytes. *Crit Care Med* 2004; 32: 1715-1721.
17. Becker KL, Nylén ES, White JC, Müller B, Snider RH. Procalcitonin and the calcitonin gene family of peptides in inflammation, infection, and sepsis: a journey from calcitonin back to its precursors. *J Clin Endocrinol Metab* 2004; 89: 1512-1525.
18. Wang H, Yin F, Shen DX, Zhang YJ, Luo YP, Liu CJ. Predictive value of procalcitonin for excluding bloodstream infection: Results of a retrospective study and utility of a rapid, quantitative test for procalcitonin. *Journal of International Medical Research* 2013; 41: 1671-1681.

19. Jensen JU, Heslet L, Jensen TH, Espersen K, Steffensen P, Tvede M. Procalcitonin increases in early identification of critically ill patients at high risk of mortality. *Crit Care Med* 2006; 34: 2596-2602.
20. Ruiz-Alvarez MJ, García-Valdecasas S, De Pablo R. Diagnostic efficacy and prognostic value of serum procalcitonin concentration in patients with suspected sepsis. *J Intensive Care Med* 2009; 24: 63-71.
21. Pettila V, Hynninen M, Takkunen O. Predictive value of procalcitonin and interleukin-6 in critically ill patients with suspected sepsis. *Intensive Care Med* 2002; 28: 1220-1225.

Correspondence to:

Sunil Kumar Nanda
Department of Biochemistry
Pondicherry Institute of Medical Sciences
Affiliated to Pondicherry University
Pondicherry