

Effect of *Centella asiatica* L. extract against ethinylestradiol induced genotoxic damage on cultured human lymphocytes

Author(s): Yasir Hasan Siddique, Tanveer Beg, Mohammad Afzal

Vol. 20, No. 2 (2009-05 - 2009-08)

Biomedical Research 2009; 20 (2): 141-147

ISSN: 0970938X

Yasir Hasan Siddique, Tanveer Beg, Mohammad Afzal

Human Genetics and Toxicology Laboratory, Section of Genetics, Department of Zoology, Faculty of Life Sciences, Aligarh Muslim University, Aligarh, India.

Abstract

In the present study the effect of *Centella asiatica* L. extract was studied against the genotoxic doses of ethinylestradiol on human lymphocytes culture. The different doses of *C. asiatica* L. extract i.e. 1.075×10^{-4} , 2.125×10^{-4} and 3.15×10^{-4} g/ml were treated separately with 5 μ M of ethinylestradiol. Similar treatments were given with 10 μ M of ethinylestradiol. The treatments result in a significant dose dependent decrease in chromosomal aberrations and sister chromatid exchanges on human lymphocyte cultures induced by 5 and 10 μ M of ethinylestradiol. The selected doses of *C. asiatica* L. extract were not genotoxic itself. Hence it is concluded that *C. asiatica* L. extract reduced the genotoxic damage during the ethinylestradiol therapy in patients and thereby reducing the chances of cancer development in humans.

Key words: *Centella asiatica* L., human lymphocytes, chromosomal aberrations, sister chromatid exchanges, natural products.

Accepted March 07 2009

Introduction

Centella asiatica L. belongs to the family Umbellifera. It is known as mandukparni or Indian Penny Wort. It is found in swampy areas of India, commonly found as a weed in crop fields and in other waste places throughout India up to an altitude of 600 m [1]. The crude extract of *C. asiatica* and the products derived from it have various pharmacological properties [2,3,4,5,6]. The crude extract of *C. asiatica* was shown to be non-toxic in normal human lymphocytes [7], and reduced the genotoxic effects of methyl methanesulphonate and cyclophosphamide in cultured human lymphocytes [8].

Conjugated estrogens, have been listed in the fourth annual report on carcinogenesis (ROC), as human carcinogens [9]. A number of and mestranol β individual steroidal estrogens including estradiol-17 were listed in the ROC as reasonably anticipated human carcinogens [9]. Steroidal estrogens are placed in a Group 1 as carcinogenic to humans, on the basis of sufficient evidence of carcinogenicity in humans [10]. Ethinylestradiol is used with various progestogens, in combined oral contraceptive formulations [11]. There are sufficient evidences for the carcinogenicity of ethinylestradiol in experimental animals [12]. It has also been reported to cause aneuploidy in Chinese Hamster DON Cells in vitro [13], aneuploidy and polyploidy both, in V79 cells in vitro Estrogens are used in the treatment of sexual and metabolic disorders and also in oral contraceptives [15]. Prolonged use of estrogens has been reported to develop various types of malignancies in human and experimental animals [15]. The genotoxic effects of estrogens/ progestins can be reduced by the use of antioxidants [16-22] and natural plant products [23-28]. The genotoxicity testing provides human a risk assessment. An increase in the frequency of chromosomal aberrations in peripheral blood lymphocytes is associated with an increased overall risk of cancer [29,30].

Any reduction in the genotoxicity gives an indication of the lesser possibility of carcinogenesis [31]. The majority of populations use traditionally natural preparations derived from the plant material for the treatment of various diseases, and for that reason it becomes necessary to assess the mutagenic potential or modulating action of plant extract in combination with other substances. In earlier study ethinylestradiol was found to be genotoxic at 5 and 10 μ M [32]. The herbal preparations can be useful in taking care of the genotoxic effects of certain drugs. The aim of the present work was to study the effect of *C. asiatica* L. extract against the genotoxic doses of ethinylestradiol.

Materials and Methods

Chemicals

Ethinylestradiol (CAS No.: 57-63-6, Sigma); RPMI 1640, Fetal calf serum, Phytohaemagglutinin-M, Antibiotic-antimycotic mixture (Gibco); Dimethyl sulphoxide, 5-Bromo-2-deoxy uridine; Colchicine (SRL, India), Giemsa stain (Merk).

Extract preparation

C. asiatica L. leaves were collected from the nursery of Forest Research Institute (FRI), Dehradun (UA) and were air dried and ground to fine powder. Extraction was performed by soaking samples (30 g of dry weight) in 300 ml of acetone for 8-10h at 40-60°C in Soxhlet's apparatus. After filtration, the excess of solvent was removed by rotatory evaporator. The extract concentrations of 1.075×10^{-4} , 2.127×10^{-4} and 3.15×10^{-4} g/ml of culture medium were established [8].

Human lymphocyte culture

Duplicate peripheral blood cultures were prepared according to Carballo, Alvarez, Boveris. [33]. Briefly, heparinized blood samples (0.5 ml), were obtained from healthy female donors and were placed in a sterile culture bottles containing 7 ml of RPMI-1640 medium, supplemented with fetal calf serum (1.5 ml), antibiotic-antimycotic mixture (1.0 ml) and phytohaemagglutinin (0.1 ml). The culture bottles were kept in an incubator at 37°C for 24 h.

Chromosomal aberration analysis

After 24 h, 5 µM of ethinylestradiol treatment (dissolved in dimethylsulphoxide, 5 µl/ml) was given separately with 1.075×10^{-4} , 2.125×10^{-4} and 3.15×10^{-4} g/ml of *Centella asiatica* L. extract. Similarly, 10 µM of ethinylestradiol was also given with the same three doses of *C. asiatica* extract. About, 0.5 ml of S9 mix (supplement with NADP) was also given with the treatments for 6 h. The cultures were washed with fresh RPMI-1640 and kept for another 48 h in an incubator at 37°C. After 47 h, 0.2 ml of colchicine (0.2 µg/ml) was added to the culture bottles. Cells were centrifuged at 800 g for 10 min. The supernatant was removed and 5 ml of prewarmed (37°C), KCl hypotonic solution (0.075 M) was added. Cells were resuspended and incubated at 37°C for 15 min. The supernatant was removed by centrifugation at 800 g for 10 min and 5 ml of chilled fixative (methanol: glacial acetic acid; 3:1) was added. The fixative was removed by centrifugation and the procedure was repeated twice. The slides were stained in 3% Giemsa solution in phosphate buffer (pH 6.8) for 15 min. Three hundred metaphases were examined for the occurrence of different types of abnormally. Criteria to classify the different types of aberrations were in accordance with recommendation of EHC46 for environmental monitoring of human population [34].

Sister chromatid exchange analysis

For sister chromatid exchange analysis, bromodeoxyuridine (10 µg/ml) was added at the beginning of the culture. After 24 h, 5 µM of ethinylestradiol (dissolved in dimethylsulphoxide, 5 µl), the treatment was given separately with 1.075×10^{-4} , 2.125×10^{-4} and 3.15×10^{-4} g/ml of *C. asiatica* extract respectively. Similar treatments with the three dosages of *C. asiatica* extract were given separately with 10 µM of ethinylestradiol. About, 0.5 ml of S9 mix (supplemented with NADP) was also given along with the treatments for 6 h. The cultures were washed with fresh RPMI 1640 and kept for another 48 h, in an incubator. Mitotic arrest was performed by adding 0.2 ml of colchicine (0.2 µg/ml). Hypotonic treatment and fixation were performed in the same way as described for the chromosomal aberration analysis. The sister chromatid exchange average was taken from an analysis of 50 metaphases during second cycle of division [35].

Statistical analysis

Student 't'-test was used for analysis of CAs and SCEs. Regression analysis was performed using Statistica Soft Inc.

Results

Ethinylestradiol induced a significant increase of abnormal metaphases as compared to the untreated at 5 and 10 μM . A significant dose dependent decrease in number of abnormal metaphase was observed when 5 and 10 μM of ethinylestradiol was treated separately, with the different doses of *C. asiatica* extract, i.e. 1.075×10^{-4} , 2.125×10^{-4} and 3.15×10^{-4} g/ml (Table 1). For sister chromatid exchange analysis, a significant increase was observed at both of the studied doses of ethinylestradiol i.e. 5 and 10 μM (Table 2). A significant decrease in sister chromatid exchanges per cell was observed when 5 and 10 μM of ethinylestradiol was treated, separately, with the different doses of *C. asiatica* extract i.e. 1.075×10^{-4} , 2.125×10^{-4} and 3.15×10^{-4} g/ml (Table 2). Regression analysis was also performed to determined the dose effect of *C. asiatica* extract on 5 and 10 μM of ethinylestradiol. For number of abnormal metaphases and sister chromatid exchanges. A decrease in the slope of linear regression lines was observed as the dose of the extract was increase in each of the treatment. For abnormal metaphases the treatment of 5 μM ($F = 29.14$; $P < 0.04$) and 10 μM ($F = 12.65$; $P < 0.05$) of ethinylestradiol, with the increase in the doses of *C. asiatica* extract results in the decrease in the slope of the linear regression lines (Figs. 1 and 2). Effect of *Centella asiatica* L. extract against ethinylestradiol genotoxicity

Table 1: Effect of *Centella asiatica* L. extract on chromosomal aberrations (CAs) induced by ethinylestradiol.

Treatments	Cells scored	Abnormal metaphases without gaps		Total structural chromosomal aberrations		
		Number	Mean \pm SE	Gaps	CTB	CSB
Ethinylestradiol (μM)						
5	300	12	4.00 \pm 1.13 ^a	9	8	4
10	300	18	6.00 \pm 1.37 ^a	11	12	6
Ethinylestradiol (μM) + CAE (g/ml)						
5+ 1.075×10^{-4}	300	8	2.67 \pm 0.93 ^b	4	6	2
5+ 2.125×10^{-4}	300	6	2.00 \pm 0.80 ^b	3	4	2
5+ 3.15×10^{-4}	300	5	1.67 \pm 0.73 ^b	2	4	1
10+ 1.075×10^{-4}	300	13	4.33 \pm 1.17 ^b	5	9	4
10+ 2.125×10^{-4}	300	10	3.33 \pm 1.03 ^b	4	6	4
10+ 3.15×10^{-4}	300	9	3.00 \pm 0.98 ^b	3	6	3
CAE (g/ml)						
1.075×10^{-4}	300	2	0.67 \pm 0.47	1	2	-
2.125×10^{-4}	300	3	1.00 \pm 0.57	2	2	1

3.15×10^{-4}	300	4	1.33 ± 0.66	2	3	1
Untreated	300	2	0.67 ± 0.47	3	2	-
Negative control (DMSO, 5 μ l/ml)	300	2	0.67 ± 0.47	1	2	-
Positive control (CP, 0.16 μ g/ml)	300	29	$9.67 \pm 1.70a$	16	18	10

CAE: Centella asiatica extract; DMSO: Dimethylsulphoxide; CP: Cyclophosphamide; CTB: Chromatid break; CSB: Chromosome break; CTE: Chromatid exchange; DIC: Dicentric; S.E.: Standard error.

Table 2: Effect of Centella asiatica L. extract on sister chromatid exchanges (SCEs) induced by ethinylestradiol.

Treatment	SCEs / Cell (Mean \pm SE)	Range
Ethinylestradiol (μ M)		
5	$6.92 \pm 0.72a$	2-8
10	$7.62 \pm 0.84a$	2-9
Ethinylestradiol (μ M) + CAE (g/ml)		
5+ 1.075×10^{-4}	$3.24 \pm 0.34b$	0-5
5+ 2.125×10^{-4}	$2.94 \pm 0.29b$	0-5
5+ 3.15×10^{-4}	$2.78 \pm 0.26b$	0-5
10+ 1.075×10^{-4}	$4.88 \pm 0.54b$	1-5
10+ 2.125×10^{-4}	$3.94 \pm 0.42b$	0-5
10+ 3.15×10^{-4}	$3.54 \pm 0.38b$	0-5
CAE (g/ml)		
1.075×10^{-4}	2.18 ± 0.18	0-5
2.125×10^{-4}	2.24 ± 0.21	0-5
3.15×10^{-4}	2.72 ± 0.23	0-5
Untreated	1.12 ± 0.12	0-5
Negative control (DMSO, 5 μ l/ml)	1.88 ± 0.16	0-5
Positive control (CP, 0.16 μ g/ml)	$12.24 \pm 0.96a$	3-13

For sister chromatid exchange analysis, the treatment of 5 μ M ($F = 35.19$; $P < 0.01$) and 10 μ M ($F = 19.69$; $P < 0.03$) of ethinylestradiol, with the increase in the dosages of *C. asiatica* extract, the decrease in slope of linear regression lines was observed (Fig. 3 and 4).




Fig. 1: Regression analysis for the dose effect of *Centella asiatica* extract on number of abnormal metaphases treated with 5 μ M of ethinylestradiol. CAE: *Centella asiatica* extract; DMSO: Dimethylsulphoxide; CP: Cyclophosphamide; S.E.: Standard error. Siddique/ Beg/Afzal




Fig. 2: Regression analysis for the dose effect of *Centella asiatica* extract on number of abnormal metaphases treated with 10 μ M of ethinylestradiol.




Fig. 3: Regression analysis for the dose effect of *Centella asiatica* extract on sister chromatid exchanges / cell treated with 5 μ M of ethinylestradiol.

Fig. 4: Regression analysis for the dose effect of *Centella asiatica* extract on sister chromatid exchanges / cell treated with 10 μM of ethinylestradiol. Effect of *Centella asiatica* L. extract against ethinylestradiol genotoxicity

Discussion

The results of the study reveal that the selected dosages of the plant extract were not genotoxic per se, but reduced the genotoxic damage of ethinylestradiol on human lymphocytes in vitro. In our earlier study, three doses of ethinylestradiol (1, 5 and 10 μM) were studied [32]. Ethinylestradiol was found to be genotoxic at 5 and 10 μM only in the presence of S9 mix supplemented with NADP. International Agency on Cancer (IARC), mainly on the basis of epidemiological studies classifies steroidal estrogens and estrogen progestins combinations among agents carcinogenic to humans (Group 1), progestins as possibly carcinogenic (Group 2) and androgenic anabolic steroids, as probably carcinogenic (Group 2A) [36]. Cancer is characterized by genetic instability, i.e. gross chromosomal abnormalities, which consist of translocation and gene amplifications. Molecular and cytogenetic evidence indicates that the induction of structural and numerical chromosome aberrations in cell play an important role in the neoplastic development of certain tumors [37]. An increase in the frequency of chromosomal aberrations in peripheral blood lymphocytes is associated with an increased overall risk of cancer [29,30]. Genetic toxicology tests are in vitro and in vivo assays designed to detect compounds that induce genetic damage directly or indirectly. Any kind of chromosomal damage may play an important role in many malignancies. The ready quantifiable nature of sister chromatid exchanges with high sensitivity for revealing toxicant-DNA interaction and the demonstrated ability of genotoxic chemicals to induce significant increase in sister chromatid exchanges in cultured cells has resulted this endpoint being used as indicator of DNA damage in blood lymphocytes of individual exposed to genotoxic carcinogens [31]. The above genotoxic endpoints are well known markers of genotoxicity and any reduction in the frequency of these genotoxic endpoints gives us an indication of the anti-genotoxicity of a particular compound [31].

At present much attention of preventive medicine research is focused on natural antioxidants, leading to isolation and identification of new biologically active molecules by the pharmaceutical industry. The crude extract of plants may contribute to total intake of plant antioxidants and be even better source of dietary antioxidants [38].

The compounds present in the extract may inhibit the promutagen bioactivation by blocking the oxidation processes or reacts with electrophilic metabolites of promutagens [39]. The verification of the possible mutagenic and/or anti-mutagenic effects of medicinal plants infusion/ extracts is another important factor in studies. Such effects have been elucidated in some plant species by using various test systems [40]. Medicinal herbs contain complex mixtures of thousand of compounds that can exert their antioxidant and free radical scavenging effect either separately or in synergistic ways [41]. Many plant products protect against xenobiotics either by inducing detoxifying enzymes or by inhibiting oxidative enzymes [42]. Detoxification systems play a major role in preventing carcinogenesis, but also an important role in preventing xenobiotic as

well as endogenous toxicity. Phytochemicals play an important role in inhibition of carcinogenesis [43]. Some plant extract may possess substances that can modulate the genotoxicity of other compounds. The data obtained in the present study suggests that the compounds present in the extract of *C. asiatica* are not mutagenic on their own or when associated with ethinylestradiol in the presence of NADP.

The protective effect observed in the present study i.e. significant reduction in the frequency of cells with chromosomal damage and sister chromatid exchanges may be due to the direct action of the compounds present in the extract on ethinylestradiol by inactivating it enzymatically or chemically. The genotoxic effects of ethinylestradiol have been attributed to the bio-activation of ethinylestradiol in the presence of S9 mix supplemented with NADP. The compounds present in the extract may also enhance the DNA repair system or DNA synthesis or even may prevent the bioactivation of certain chemicals [44]. Our earlier reports on synthetic progestins showed that they are genotoxic in *in vitro* [45,46,47,48] and *in vivo*. A reduction in the frequency of SCE and chromosomal aberrations in the test system thereby indicates the possibility of reducing the chances of carcinogenesis during the ethinylestradiol therapy in patients.

Conclusion

The extract of *Centella asiatica* L. is potent enough to reduce the genotoxic damage of ethinylestradiol. The results of the present study suggest that the plant extract can form the basis of herbal medicine as it has antioxidants that are responsible for the reduction of the genotoxic damage of the drugs. The prolonged use of ethinyl estradiol has been reported to induce cancer. Any reduction in the frequency of CAs and SCEs thereby gives an indication of little chances of developing cancer. The extract of *Centella asiatica* reduces the frequency CAs and SCEs induced by ethinyl estradiol.

Acknowledgements

Thanks are due to the DST, New Delhi for awarding the project under SERC-Fast Track Scheme No: SR/FT/LS-003/2007 to the author (YHS) and to the Chairman, Department of Zoology, Aligarh Muslim University, Aligarh, India, for laboratory facilities.

References

1. Dastur JF. Medicinal plants of India and Pakistan. D.B. Taraporevala Sons & Co., Bombay, 1962.
2. Dutta T, Basu UP. Crude extract of *Centella asiatica* and products derived from its glycosides as oral antifertility agents. *Indian J Exp Biol* 1968; 6: 181-182.
3. Guo JS, Cheng CL, Koo MW. Inhibitory effects of *Centella asiatica* water extract and asiaticoside on inducible nitric oxide synthase during gastric ulcer healing in rats. *Planta Medica* 2004; 70: 1150-1154.
4. Punuree K, Wild CP, Kasinrerk W, Vinitketkumnuen U. Immunomodulatory activities of *Centella asiatica* and *Rhimachanthus nasutus* extracts. *Asian Pacific J Can Prev* 2005; 6: 396-400.

5. Inamdar PK, Yeole KD, Cibogare AB, Desouza NJ. Determination of biologically active constituents in *Centella asiatica*. *J Chromatography* 1996; 742: 127-130.
6. Yoshida M, Fuchigami M, Nago T, Okabe H, Matsunaga K, Takata J, Karube J, Tsuchihashi R, Kinjo J, Mihashi K and Fujioka T. Antiproliferative constituents from Umbelliferae plants VII. Active triterpene and rosmarinic acid from *Centella asiatica*. *Biol Pharmacol Bull* 2005; 28: 173-175.
7. Babu TD, Kuttar G, Padikkla J. Cytotoxic and antitumor properties of certain taxa of umbelliferae with special reference to *Centella asiatica* (L.) urban. *J Ethnopharmacol* 1995; 48: 53-57.
8. Siddique YH, Ara G, Beg T, Faisal M, Ahmad M, Afzal M. Protective role of *Centella asiatica* L. extract against methylmethane sulphonate and cyclophosphamide induced genotoxic damage in cultured human lymphocytes. In: *Phytopharmacology and Therapeutic Values I*, Studium Press LLC, USA. *Recent Prog Med Plants* 2007; 19: 369-381.
9. ROC Fourth Report on Carcinogenesis, US DHHS, National Toxicology Program 1985.
10. IARC. Overall evaluations of carcinogenicity an updating of IARC monographs, Vol. 1-42, IARC Monographs on the Evaluation of Carcinogenic Risks to humans (Suppl. 7). International Agency for Research on Cancer, Lyon, France, 1987; p. 280.
11. Schwend TH, Lippman JS. Comparative review of recently introduced oral contraceptives containing norgestimate, desogestrel and gestodene and older oral contraceptives. In: Paulik EJ (Ed.) *Estrogens, Progestins and their antagonists*, Birkhauser, Boston, 1996; pp. 273-296.
12. IARC. Sex hormones (II), IARC Monograph on the Evaluations of carcinogenic risk of chemicals to humans, Vol. 21, International Agency for Research on Cancer, Lyon, France 1979.
13. Wheeler WJ, Cherry LM, Downs T, Hsu TC. Mitotic inhibition and aneuploidy induction by naturally occurring and synthetic estrogens in Chinese Hamster Cells in vitro. *Mutat Res* 1986; 171: 31-41.
14. Sato Y, Sakakibara Y, Oda T, Aizu-Yokota E, Ichinoseki K. Effect of estradiol and ethinylestradiol on microtubule distribution in Chinese Hamster V79 Cells. *Chem Pharmacol Bull* 1992; 40: 182-184.
15. IARC. Post menopausal oestrogen therapy. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans, Vol. 72, International Agency for Research on Cancer, Lyon, France, 1999; p. 399.
16. Siddique YH, Ara G, Beg T, Afzal M. Additive action of vitamin C and E against norgestrel induced genotoxicity. *Biomed Res* 2007; 18: 155-160.
17. Siddique YH, Beg T, Afzal M. Antigenotoxic effects of ascorbic acid against megestrol acetate induced genotoxicity in mice. *Hum Exp Toxicol* 2005; 24: 121-127.
18. Siddique YH, Ara G, Beg T, Afzal M. Effect of vitamin C on cyproterone acetate induced genotoxic damage in mice. *Res J Biol Sci* 2006 1: 69-73.
19. Siddique YH, Beg T, Afzal M. Anticlastogenic effects of ascorbic acid against the genotoxic damage induced by norethynodrel. *Adv Environ Biol* 2007; 1: 27-32.

20. Siddique YH, Afzal M. Protective role of allicin and L. ascorbic acid against the genotoxic damage induced by chlormadinone acetate in cultured human lymphocytes. *Indian J Exp Biol* 2005; 49: 769-772.
21. Siddique YH, Beg T, Ara G, Gupta J, Afzal M. Antigenotoxic effect induced genotoxic damage in cultured β of allicin against estradiol-17 mammalian cells. *Natural Products Research* (in press)
22. Siddique YH, Beg T., Afzal M. Antigenotoxic effect of ascorbic acid against cyproterone acetate induced genotoxicity in cultured mammalian cells. In: *Recent Trends in Toxicology*, YH. Siddique (Eds), Transworld Research Network, Trivandpuram, Kerala, India. 2008 pp. 85-94.
23. Siddique YH, Ara G, Beg T, Afzal M. Antigenotoxic effect of *Ocimum sanctum* L. extract against cyproterone acetate induced genotoxic damage in cultured mammalian cells. *Acta Biol Hungarica* 2007; 58: 397-407.
24. Siddique YH, Ara G, Beg T, Afzal M. Antigenotoxic effect of nordihydroguaiaretic acid (NDGA) against chlormadinone acetate induced genotoxic damage in mice bone marrow cells. *J Nat Med* 2008; 62: 52-56.
25. Siddique YH, Beg T, Afzal M. Protective effect of nordihydroguaiaretic acid (NADGA) against norgestrel induced genotoxic damage. *Toxicol in vitro* 2006; 20: 227-233.
26. Siddique YH., Ara G., Beg T., Faisal M., Ahmad M., Afzal M. Antigenotoxic role of *Centella asiatica* L. extract against cyproterone acetate induced genotoxic damage in cultured human lymphocytes. *Toxicol invitro*. 2008. 22: 10-17.
27. Siddique YH., Ara G., Beg T, Afzal M. Possible modulating action of plant infusion of *Ocimum sanctum* L. on chromosomal aberrations and sister chromatid exchanges induced by chlormadinone acetate in human lymphocytes invitro. *J. Environ. Biol.* 2008; 845-848.
28. Siddique YH., Ara G, Beg T, Afzal M. Protective role of nordihydroguaiaretic acid (NDGA) against the genotoxic damage induced by ethinodiol diacetate in human lymphocytes invitro. *J Environ Biol.* 2007; 28: 279-282.
29. Hagmar L, Bonassi S, Stromberg U, Brogger A, Knudson JE, Norppa H, Reuterwall C. Chromosomal aberrations in human lymphocytes predicts human cancer: A report from the European study group on cytogenetic biomarkers and health (ESCH). *Cancer Res* 1998; 58: 4117-4121.
30. Hagmar L, Brogger A, Hansteen IL, Heim S, Hogstedt B, Knudsen L, Lambert B, Linnainmaa K, Mitelman F, Nordenson I, Reuterwall C, Salomaa S, Skerfving S, Sorsa M. Cancer risk in humans predicted by increased level of chromosomal aberration in human lymphocytes. Nordic study group on the health risk of chromosome damage. *Cancer Res* 1994; 54: 2919-2922.
31. Albertini RJ, Anderson D, Douglas GR, Hagmar L, Heminki K, Merelo F, Natrajan AT, Norppa H, Shuker DEG, Tice R, Walters MD, Aitio A. IPCS guidelines for the monitoring of genotoxic effects of carcinogens in humans. *Mutat Res* 2000; 463: 119-172.
32. Siddique YH, Beg T, Afzal M. Genotoxic potential of ethinylestradiol in cultured mammalian cells. *Chem Biol Interact* 2005; 151: 133-141.
33. Carballo MA, Alvarez S, Boveris S. Cellular stress by light and Rose Bengal in human lymphocytes. *Mutat Res* 1993; 288: 215-222.

34. IPCS. Internatinal Programme on Chemical Safety: Environment Health Criteria 46. Guidelines for the study of genetic effects in human populations, WHO Geneva, 1985; pp. 25-54.
35. Perry P, Wolff S. New Giemsa methods for differential staining of sister chromatids. *Nature* 1974; 261: 156-158.
36. Martelli A, Mattioli F, Angiola M, Reimann R, Brambilla G. Species, sex and inter-individual differences in DNA repair induced by nine sex steroids in primary culture of rat and human hepatocytes. *Mutat Res* 2003; 536: 69-78.
37. Yunis, JJ. The chromosomal basis of human neoplasia. *Science* 1983; 221: 227-236.
38. Zamorano-Ponce E, Morales C, Ramos D, Sepulveda C, Cares S, Rivera P, Fernandez J, Carballo MA. Antigenotoxic effect of Aloysia triphylla infusion against acrylamide induced DNA damage as shown by the comet assay technique. *Mutat Res* 2006; 603: 145-150.
39. Kaya, B Antigenotoxic effect of ascorbic acid on mutagenic dose of three alkylating agents. *Turkish J Biol* 2003; 27: 241-246.
40. Roncada T, Vicentini VEP, Mantovani MS. Possible modulating actions of plant extracts on the chromosome breaking activity of MMC and Ara C in human lymphocytes in vitro. *Toxicol in vitro* 2004; 18: 617622.
41. Romero-Jimenez M, Sanchez JC, Analla M, Serrano AM, Moraga AA. Genotoxicity and antigenotoxicity of some traditional medicinal herbs. *Mutat Res* 2005; 585: 147-155.
42. Morse MA, Stoner GD. Cancer chemoprevention: principles and prospects. *Carcinogenesis* 1993; 14: 1737-1746.
43. Blaylock, RL. A review of conventional cancer prevention and treatment and the adjunctive use of nutraceutical supplements and antioxidants: Is there a danger or a significant benefit? *J Am Nutr Assoc* 2000; 3: 75-95.
44. Kuroda Y, Jain AK, Tezuka H, Kada T. Antimutagenicity in cultured mammalian cells. *Mutat Res* 1992; 267: 201-209.
45. Siddique YH, Ara G., Beg T, Afzal M. Genotoxic potential of medroxyprogesterone acetate in cultured human peripheral blood lymphocytes. *Life Sci.* 2006; 80: 212-218.
46. Siddique YH, Afzal M. Genotoxic potential of cyproterone acetate of possible role of reactive oxygen species. *Toxicol in vitro.* 2005; 19: 63-68.
47. Siddique YH, Afzal M. Evaluation of genotoxic potential of norethynondrel in human lymphocytes in vitro. *J. Environ Biol* 2005; 26: 387-392.
48. Siddique YH, Afzal M. Evaluation of genotoxic potential of ethynodiol diacetate in human lymphocytes in vitro. *Curr.Sci.* 2004; 86: 1161-1165.
49. Siddique YH, Afzal M. Evaluation of genotoxic potential of synthetic progestin chlormadinone acetate. *Toxicol Lett* 2004; 15: 221-225.
50. Siddique YH, Afzal M. In vivo evaluation of sister chromatid exchanges (SCEs) and chromosomal aberrations (CAs) by lynestrenol. *Indian J Exp Boil* 2005; 43: 291-293.

Correspondence:

Yasir Hasan Siddique

Human Genetics and Toxicology Laboratory, Section of Genetics
Department of Zoology, Faculty of Life Sciences, Aligarh Muslim University
Aligarh 202002 (U.P.), India