

Effect of comprehensive nursing intervention on therapy and life quality of patients with breast cancer.

Xia Kang^{1*}, Yu Fang², Yu Kong²

¹Department of Imaging, Zhengzhou People's Hospital, Zhengzhou, PR China

²General ICU, the First Affiliated Hospital of Zhengzhou University, Zhengzhou, PR China

Abstract

Objective: Many cancer patients suffered from mental disorders. This research was to investigate the effect of comprehensive nursing intervention on patients with breast cancer.

Method: Patients were accepted routine nursing in control group (group C), and comprehensive nursing intervention in intervention group (group I). At the end of the operation and in the following two months after operation, the levels of depression, anxiety, life quality, and sleep quality were evaluated, and the expressions of Lactic Dehydrogenase (LDH), α -fetoprotein (AFP) and Carcino-Embryonic Antigen (CEA) were also detected. The rates of recurrence and metastasis were assessed during 3-year follow-up.

Results: The levels of depression and anxiety of patients in group I were much higher than group C ($P<0.05$). Life and sleep qualities in both groups were improved, while the quality in group I was significantly better than group C ($P<0.05$). The expressions of LDH, CEA and AFP of patients in both groups were reduced, while the reduction in group I was more significant than group C ($P<0.05$). After 3-year follow-up, the recurrence and metastasis rates in group I were significantly lower than group C ($P=0.044$, $P<0.05$).

Conclusions: Comprehensive nursing intervention contributed to the improvement of life quality, sleep quality and mental health, and was helpful to improve the therapeutic effect of breast cancer.

Keywords: Comprehensive nursing interventions, Breast cancer, Depression, Anxiety, Life quality, Sleep quality.

Accepted on September 28, 2017

Introduction

It is time to no longer consider breast cancer as a single disease. Different subtypes have differences in risk factors [1,2], natural histories [3], responses to systemic and local therapies [4,5], and genomic atlas.

The potentially informed treatment decision should be made according to these characters [6]. Otherwise, researches showed that many cancer patients would suffer from some mental disorders, such as depression and anxiety.

Mental trouble in cancer patients could bring significantly financial and emotional burden, which would result in the high utilization of health services, low quality of life and the increase of mortality [7]. Detailed treatment decision about breast cancer should be made in the consideration of disease extent, host factors, patient preferences, and social and economic constraints [8]. And the nursing serves should pay more attention to the mental health.

In our research, we implemented the comprehensive nursing intervention for patients with breast cancer to evaluate its influence.

Methods

Patients

200 patients with breast cancer, who were confirmed by our hospital from January 2010 to December 2015, were randomly divided into two groups to compare the clinical treatment effect of different nursing care. There were respectively 100 patients in control group (group C) and intervention group (group I) in which all patients were younger than 40 years old. There was no significant difference between two groups in age, gender, disease course, cancer grade and educational background.

The following were the inclusion criteria of patients: 1) Patients were 18-40 y old. 2) Breast cancer was in I to III grades. 3) Patients had the ability of communication and reading. 4) Patients did not have the history and family history of psychiatric illness. 5) Patients were never diagnosed as other cancers. 6) The patient could undergo biopsy or surgery in a primary tumor site, which was suspected or proven as breast cancer. 7) Patients were treated with the operation of breast cancer and clinical postoperative chemotherapy. 8) Patients were without other severe diseases. 9) The patient was willing

to participate in this research after informed the content. Patients had accepted radiation treatment would be excluded. And patients with other cancers or breast cancer recurrence were also not involved.

Nursing management

Patients in group C accepted routine nursing for breast cancer which included the general nursing, health propaganda and education in and out of hospital. In addition to the routine nursing, patients in group I were implemented comprehensive nursing intervention, including psychological nursing, behavior and life nursing intervention, social intervention, health education and music relaxation.

Evaluation of life quality

The life quality of all patients was evaluated with EORTC QLQ-C30 questionnaire (European Organization for Research and Treatment of Cancer quality of life questionnaire) in 15 fields [9]. This evaluation was managed after the breast cancer operation, and when reexamined after 1- and 2-month nursing.

Evaluation of depression and anxiety

The Self-Rating Depression Scale (SDS) [10] was used for the assessment of depression. And Self-Rating Anxiety Scale (SAS) [11] was applied to the evaluation of anxiety. The following was the standard of evaluation (Table 1).

We counted up the number of patients with depression or anxiety in groups C and I. After 1- and 2-month nursing, these would be managed when re-examined.

Evaluation of sleep quality

Sleep quality was assessed by using Pittsburgh Sleep Quality Index (PSQI) [12], which was to produce the global sleep quality index score in a 19-item scale on the basis of seven component indexes: sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbance, use of sleep medications and daytime dysfunction. This assessment also would be implemented post operation and when reexamined after 1- and 2-month nursing.

Evaluation of therapeutic effect

The Lactic Dehydrogenase (LDH), α -fetoprotein (AFP) and Carcino-Embryonic Antigen (CEA) were detected post operation. These indexes would be detected again when re-examined after 1- and 2-month nursing.

Statistical analysis

Statistical analysis was performed with the SPSS, version 21.0. All measurement data were presented as mean \pm standard deviation and analysed with T test. P value with less than 0.05 indicated a statistically significant difference and was adjusted for multiple comparisons.

Results

Basic information

200 patients participated in our research. There were 100 patients in each group. The statistical analysis results showed that patients in two groups were not significantly different in age, marital status, educational background and pathological stage ($P > 0.05$) (Table 2).

Evaluation of life quality

With the EORTC QLQ-C30 questionnaire, we detected the life quality of patient in two groups. Just after the breast cancer operation, the scores of patients in two groups were respectively 40.7 ± 12.1 and 39.3 ± 14.6 . There was no significant difference ($P = 0.65$, $P > 0.05$). Then after two-month nursing, life qualities of all patients were increased. However, the life quality of group I was much better than group C, with a statistically significant difference ($P < 0.05$) (Table 3).

Evaluation of Sleep quality

PSQI scale assessed the quality of sleep in 7 components. There was no significant difference between two groups after breast cancer operation ($P < 0.05$). However, after two-month comprehensive nursing intervention, the sleep quality of group I was much better than group C. And the difference was statistically significant ($P < 0.05$). Moreover, this difference reflected in each component (Table 4).

Evaluation of depression and anxiety

With SDS and SAS scale, we evaluated the levels of depression and anxiety for patients with breast cancer. In group I, there were 38 (38%) depression patients and 35 (35%) anxiety patients, while 37 (37%) depression patients and 33 (33%) anxiety patients in group C. Though there was no difference between two groups just after the surgery, a remarkable reduction appeared after two month careful nursing ($P < 0.05$). Moreover, the difference between two groups was more and more remarkable and statistically significant ($P < 0.05$). The condition of depression and anxiety of patients in group I were better than them in group C ($P < 0.05$) (Table 5).

Therapeutic results

After the operation of breast cancer and postoperative chemotherapy, patients in two groups were managed with different nursing model. We detected the expression levels of tumor markers, LDH, AFP and CEA. The expressions of these proteins were reduced both in groups I and C. But the reduction in group I was more significant than group C (Figure 1). As shown in Figure 1, after 1 or 2 month treatment and nurse, the expressions of these proteins of patients in group I were much lower than those in group C. With repeated measures analysis of variance, the statistical result showed that this difference was significant ($P < 0.05$). So, there was a better

therapeutic effect of patients in group I who accept comprehensive nursing intervention.




Figure 1. Expressions of LDH, AFP and CEA. The expressions of these proteins were reduced for patients in two groups. And in group I, the levels of these proteins were much lower than them of group C with a statistical significant ($P < 0.05$).

Recurrence of breast cancer

After 3 years follow-up, in group I, 7 cases presented with recurrent tumors, and 9 cases with tumor metastasis. Compared with group C which had 13 cases with tumor recurrence and 15 cases with tumor metastasis, the recurrence and metastasis rates in group I were significantly lower ($P = 0.044$, $P < 0.05$) (Table 6).

Table 1. The standard of depression and anxiety.

Scale	Standard	Grade
Depression (SDS)	<40	No
	40-47	Mild depression
	48-56	Moderate depression
	>56	Serious depression
Anxiety (SAS)	<44	No
	45-59	Mild-to-moderate anxiety
	60-74	Severe Anxiety
	>75	Extremely severe anxiety

Table 2. The basic information of patients in two groups.

Characteristic	Group I (n=100)	Group C (n=100)	P
Age, mean (y)*	36.9 ± 8.3	35.2 ± 7.6	0.133
Marital status (Married/other)#	92/8	90/10	0.621
Educational background (bachelor degree or above/ high school/middle school)§	39/36/25	31/39/30	0.21
Pathological stage§	34/42/14/10	30/49/11/10	0.765

Note: *Statistically analysed with t-test; #Analysed with chi-square test; §Analysed with rank sum test.

Table 3. The life quality of patients. In this table, the rows “1 month” and “2 month” respectively represent 1 month and 2 month after operation.

Index	Group	Postoperation	1 month	2 month
Life quality	I	40.7 ± 12.1	49.9 ± 14.3	58.6 ± 16.4
(EORTC QLQ-C30) ^{ab}	C	39.3 ± 14.6	42.1 ± 11.6	49.7 ± 13.5

Note: Repeated measures analysis of variance, life quality: $F(\text{management}) = 24.250$, $P < 0.05$; $F(\text{time}) = 16.012$, $P < 0.05$. LSD was used for the comparison among groups. ^athere is a difference between postoperation and 1 month or 2 month. ^bthere is a difference between 1 month and 2 month after surgery ($P < 0.05$).

Table 4. The sleep quality of patients.

Index	Group	Postoperation	1 month	2 month
Sleep quality	I	11.3 ± 2.6	10.2 ± 2.1	9.4 ± 1.8
(PSQI) ^{ab}	C	10.9 ± 2.5	12.6 ± 2.4	14.3 ± 2.6

Note: Repeated measures analysis of variance, sleep quality: $F(\text{management}) = 89.366$, $P < 0.05$; $F(\text{time}) = 23.491$, $P < 0.05$. LSD was used for the comparison among groups. ^athere is a difference between postoperation and 1 month or 2 month. ^bthere is a difference between 1 month and 2 month after surgery ($P < 0.05$).

Table 5. The depression and anxiety of two groups.

Index	Group	Postoperation	1 month	2 month
Depression (SDS) ^{ab}	I	47.3 ± 5.5	33.2 ± 4.6	29.9 ± 3.6
	C	48.1 ± 5.8	40.6 ± 5.1	36.6 ± 4.5
Anxiety (SAS) ^{ab}	I	40.5 ± 5.7	28.9 ± 3.7	25.6 ± 3.5
	C	41.2 ± 6.4	35.7 ± 4.8	32.1 ± 3.9

Note: Repeated measures analysis of variance. Depression: $F(\text{management}) = 21.126$, $P < 0.05$; $F(\text{time}) = 47.358$, $P < 0.05$; Anxiety, $F(\text{management}) = 21.126$, $P < 0.05$; $F(\text{time}) = 8.765$, $P < 0.05$. LSD was used for the comparison among groups. ^athere is a difference between postoperation and 1 month or 2 month. ^bthere is a difference between 1 month and 2 month after surgery ($P < 0.05$).

Table 6. The tumor recurrence and metastasis of two groups with 3 years follow-up.

Group	Recurrence	Metastasis	Rate	P
I (n=100)	7	9	16%	0.044
C (n=100)	13	15	28%	

Note: Chi-square test, $P < 0.05$.

Discussion

Breast is the secondary sexual characteristics of females whose function and beauty is very important [13]. The disease about breast will lead to severe psychology disorder. In our research, the patients were young whose ages were younger than 40 years. These young patients with breast cancer had high levels of depression and anxiety and low life quality and sleep quality. It had been reported that the sleep quality of younger patients with breast cancer was better than elders. However, some components of life quality were worse than older patients [14]. And some researchers had shown that the depression and anxiety levels of younger patients were higher than old persons

[15]. So, for younger patients with breast cancer, we should pay more attention to their mental troubles. The nursing model should be improved. In our research, we implemented a comprehensive nursing intervention which could preferably improve psychological status, life and sleep qualities of young patients. After two month careful nursing intervention, the evaluation of life, sleep, depression and anxiety for patients in group I was much better than patients in group C.

Moreover, LDH, CEA and AFP were important biomarkers of tumor which is a growing role in all aspect of cancer care from screening to follow-up after treatment [16]. The detection of LDH, CEA and AFP showed that the therapeutic effect of patients in group I was better than group C. And tumor recurrence and metastasis rates were lower for patients implemented with comprehensive nursing intervention. This demonstrated that well nursing intervention could exert a positive influence to the therapeutic effect. This might be because of the release of emotion and reduction of depression and anxiety. High level of emotional distress would produce a significant influence to the treatment of cancer [17]. A meta-analysis showed that patients suffering from clinical depression were three times more likely to tend to unsuccessful oncological treatment [18]. So, the mental health is important in the clinical treatment of cancers.

Moreover, nursing intervention could reduce the anxiety about economic burden induced by disease treatment. Then patients can receive a better therapy without any worries. It will be necessary to concentrate more energies on the mental health in future treatment of breast cancer.

References

1. Millikan RC, Newman B, Tse CK, Moorman PG, Conway K, Dressler LG, Smith LV, Labbok MH, Geradts J, Bensen JT, Jackson S, Nyante S, Livasy C, Carey L, Earp HS and Perou CM. Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat* 2008; 109: 123-139.
2. Phipps AI, Chlebowski RT, Prentice R, McTiernan A, Stefanick ML, Wactawski-Wende J, Kuller LH, Adams-Campbell LL, Lane D, Vitolins M, Kabat GC, Rohan TE and Li CI. Body size, physical activity, and risk of triple-negative and estrogen receptor-positive breast cancer. *Cancer Epidemiol Biomarkers Prev* 2011; 20: 454-463.
3. Phipps AI, Buist DS, Malone KE, Barlow WE, Porter PL, Kerlikowske K and Li CI. Reproductive history and risk of three breast cancer subtypes defined by three biomarkers. *Cancer Causes Control* 2011; 22: 399-405.
4. Tang G, Shak S, Paik S, Anderson SJ, Costantino JP, Geyer CE, Jr., Mamounas EP, Wickerham DL and Wolmark N. Comparison of the prognostic and predictive utilities of the 21-gene Recurrence Score assay and Adjuvant! for women with node-negative, ER-positive breast cancer: results from NSABP B-14 and NSABP B-20. *Breast Cancer Res Treat* 2011; 127: 133-142.
5. Aebi S, Sun Z, Braun D, Price KN, Castiglione-Gertsch M, Rabaglio M, Gelber RD, Crivellari D, Lindtner J, Snyder R, Karlsson P, Simoncini E, Gusterson BA, Viale G, Regan MM, Coates AS and Goldhirsch A. Differential efficacy of three cycles of CMF followed by tamoxifen in patients with ER-positive and ER-negative tumors: long-term follow up on IBCSG Trial IX. *Ann Oncol* 2011; 22: 1981-1987.
6. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. *Nature* 2012; 490: 61-70.
7. Satin JR, Linden W, Phillips MJ. Depression as a predictor of disease progression and mortality in cancer patients: a meta-analysis. *Cancer* 2009; 115: 5349-5361.
8. Goldhirsch A, Winer EP, Coates AS, Gelber RD, Piccart-Gebhart M, Thurlimann B and Senn HJ. Personalizing the treatment of women with early breast cancer: highlights of the St Gallen International Expert Consensus on the Primary Therapy of Early Breast Cancer 2013. *Ann Oncol* 2013; 24: 2206-2223.
9. Costa DS, Aaronson NK, Fayers PM, Pallant JF, Velikova G and King MT. Testing the measurement invariance of the EORTC QLQ-C30 across primary cancer sites using multi-group confirmatory factor analysis. *Qual Life Res* 2015; 24: 125-133.
10. Zung WW. A self-rating depression scale. *Arch Gen Psychiatry* 1965; 12: 63-70.
11. Zung WW. The measurement of affects: depression and anxiety. *Mod Probl Pharmacopsychiatry* 1974; 7: 170-188.
12. Otte JL, Rand KL, Carpenter JS, Russell KM and Champion VL. Factor analysis of the Pittsburgh Sleep Quality Index in breast cancer survivors. *J Pain Symptom Manage* 2013; 45: 620-627.
13. Ellis H. *Man and woman: a study of secondary and tertiary sexual characters*. Elsevier 2013.
14. Savard J, Simard S, Hervouet S, Ivers H, Lacombe L and Fradet Y. Insomnia in men treated with radical prostatectomy for prostate cancer. *Psychooncology* 2005; 14: 147-156.
15. Repetto L, Venturino A, Vercelli M, Gianni W, Biancardi V, Casella C, Granetto C, Parodi S, Rosso R and Marigliano V. Performance status and comorbidity in elderly cancer patients compared with young patients with neoplasia and elderly patients without neoplastic conditions. *Cancer* 1998; 82: 760-765.
16. Sharma S. Tumor markers in clinical practice: general principles and guidelines. *Indian J Med Paediatr Oncol* 2009; 30: 1-8.
17. Singer S, Danker H, Briest S, Dietrich A, Dietz A, Einenkel J, Papsdorf K, Lordick F, Meixensberger J, Mossner J, Niederwieser D, Prietzel T, Schiefke F, Stolzenburg JU, Wirtz H and Kersting A. Effect of a structured psycho-oncological screening and treatment model on mental health in cancer patients (STEPPED CARE): study protocol for a cluster randomized controlled trial. *Trials* 2014; 15: 482.
18. DiMatteo MR, Giordani PJ, Lepper HS, Croghan TW. Patient adherence and medical treatment outcomes: a meta-analysis. *Med Care* 2002; 40: 794-811.

***Correspondence to**

Xia Kang

Department of Imaging

Zhengzhou People's Hospital

PR China