

Effects of recombinant human interferon $\alpha 1b$ on *Bcl-2* and *c-myc* expression in chronic myeloid leukemia and correlation with clinical treatment efficacy.

Yuan He, Pengxiang Guo*, Xiaoli Du, Xue Fu

Department of Hematology, Guizhou Provincial People's Hospital, Guiyang, Guizhou, PR China

Abstract

Objective: Recombinant human interferon $\alpha 1b$ is a clinical drug in treating certain viral diseases or malignant tumors, and is also widely used in treating leukemia, although its anti-tumor mechanism remains unclear. This study aimed to investigate the effect of recombinant human interferon $\alpha 1b$ on expression of peripheral anti-tumor genes *Bcl-2* and *c-myc*, and its correlation with clinical treatment efficacy.

Patients and methods: A total of 95 chronic myeloid leukemia patients were recruited. Among all patients 46 received recombinant human interferon $\alpha 1b$ and hydroxyurea, and 49 patients received hydroxyurea treatment only. Fluorescent qRT-PCR was used to measure mRNA expression of *Bcl-2* and *c-myc* in peripheral blood of chronic myeloid leukemia patients. Treatment efficacy was evaluated from the aspect of hematology, molecular and cellular genetics. SPSS software was used to analyse the correlation between *Bcl-2* and *c-myc* mRNA expression and clinical treatment efficacy.

Results: With elongated treatment time, *Bcl-2* expression was significantly elevated after 6 w combined treatment, and *c-myc* expression was increased after 2 w combined treatment. Single hydroxyurea treatment group showed elevated levels after 6 w. Clinical treatment efficiency was positively correlated with *Bcl-2* mRNA expression ($r=0.541\sim 0.623$, $p<0.01$), and *c-myc* mRNA expression ($r=0.645\sim 0.612$, $p<0.01$).

Conclusions: Recombinant human interferon $\alpha 1b$ plays an important role in chronic myeloid leukemia treatment *via* facilitating *Bcl-2* and *c-myc* expression.

Keywords: Recombinant human interferon $\alpha 1b$, Chronic myeloid leukemia, *Bcl-2*, *c-myc*.

Accepted on July 11, 2017

Introduction

Leukemia is a type of malignant disease caused by abnormality of Hematological Stem Cells (HSCs), as colony cells lose differentiation or maturation ability for arresting at certain stage during development. Chronic Myeloid Leukemia (CML) is a hematological malignant tumor that severely threatens people's health, and is also an acquired malignant clonal disease of HSCs with 1~2 per 100,000 incidence, occupying about 15% of newly diagnosed leukemia [1]. CML is frequently featured as Ph chromosome containing *BCR-ABL1* fusion gene [2]. Pathogenesis of CML is still unclear and occupies about 3% of malignant tumors.

CML is an acquired clonal disease derived from HSCs, and is manifested as myeloid cell lineage proliferation, peripheral leukocyte hyperplasia and spleen/liver enlargement [3], accompanying with Ph chromosome and *BCR-ABL* fusion gene in bone marrow cells [4]. Slow progression of CML disease makes insidious symptoms at early stage [5]. With disease progression, leukemia can impair normal hematological function of bone marrow and cause multiple related symptoms including anemia, refractory infection and hemorrhage. Current

treatments for CML mainly focus on the chronic stage at early phase per disease pathogenesis mechanism. Common drugs for CML include imatinib, hydroxyurea, cytarabine and interferon [6,7]. By molecular target treatment and HSCs transplantation, drugs are also administrated to inhibit elevation of leukocytes. Before onset of molecular targeted drugs, interferon is the first choice of CML treatment. Recombinant human interferon α can obtain persistent cytogenetic remission of CML, thus elongating lifespan [8]. The treatment of CML using interferon has been accepted widely. Currently used interferon drugs include $\alpha 2a$ and $\alpha 2b$ interferon [9,10]. However, the treatment efficacy of interferon $\alpha 1b$ on CML has not been clearly illustrated yet.

Bcl-2 gene was firstly separated from non-Hodgkin lymphoma, and is an important tumor suppressor gene playing important roles in suppressing cell apoptosis [11]. Previous study showed that most anti-tumor drugs could inhibit leukemia cell proliferation, induce cell apoptosis and suppress *Bcl-2* expression [12]. *C-myc* gene participates in cell proliferation process, and facilitates cell apoptosis [13]. Tumorigenesis is the result of various genes and external factors. The single expression of *Bcl-2* cannot induce tumor, but only with co-

expression with oncogene *c-myc* to induce tumorigenesis [14]. To further illustrate the important mechanism of interferon $\alpha 1b$ in tumor treatment, this study investigated the effect of $\alpha 1b$ interferon on *Bcl-2* and *c-myc* expression.

Materials and Methods

Clinical information

A total of 95 CML patients admitted in hematology department of our hospital from May 2012 to October 2016 were recruited, including 57 males and 38 females. A retrospective control study was performed. 46 patients received combined treatment using both recombinant human interferon $\alpha 1b$ and hydroxyurea, and 49 patients received hydroxyurea treatment only. Basic information including age and gender distribution between two treatment groups was analysed by statistical methods.

This study has been pre-approved by the ethical committee of Guizhou Provincial People's Hospital. All subjects have signed the consent forms before recruitment in this study.

Diagnostic criteria

Inclusive criteria: (1) Anemia, uncomfortable in spleen area, hemorrhage, fatigue, body mass loss and low fever with elevated metabolic rate. (2) Leukocyte count $>50 \times 10^9/L$, with occasionally higher than $500 \times 10^9/L$, hemoglobin <110 g/L indicating normochromic anemia, plus higher platelet at $1000 \times 10^9/L$. Blood smear showed granulocytes at various maturation stages, with predominant distribution at middle and late stage of promyelocytes, as less than 5% of primordial cells and less than 10% of primary granulocyte and early promyelocytes, increased count of basophilic and acidophilic granulocytes, and minor amounts of enucleated erythrocytes. (3) Fever, spleen swelling, bone pain, hemorrhage, extra-bone marrow edema infiltration with unknown reasons at acute phase. Meanwhile, a total of 38 healthy volunteers were recruited as the control group.

Treatment methods

Skin test was performed before using recombinant human interferon $\alpha 1b$. Subcutaneous or intramuscular injection was performed using $\alpha 1b$ interferon (3000000 IU/d) (Kexing Biotech, China). Based on leukocyte count, hydroxyurea (Huarun Pharm, China) was orally applied (1-3 g/d) after 15~30 d of $\alpha 1b$ interferon treatment, accompanied with leukocyte count. Drugs were stopped or decreased when leukocyte number returned to normal level.

mRNA extraction and mRNA expression level assay

Peripheral blood samples were collected and centrifuged to obtain serum. Trizol method was used to extract tissue RNA, whose content and purity were measured by spectrometry. RNA solution after quantification was analysed for integrity using 1% agarose gel electrophoresis. 1 μ g RNA was used in reverse transcription of cDNA, following the manual

instruction of real time fluorescent quantitative RT-PCR kit (TaKaRa, Japan). cDNA after reverse transcription was added into PCR system under following conditions: 95°C denature for 5 min, followed by 40 cycles each containing 95°C 15 s and 60°C 60 s. Amplification was performed in ViiA7 fluorescent quantitative PCR cycler. Each sample was performed in triplicates. Primers were synthesized by Jierui Biotech (China). Primer sequences were shown in Table 1.

Observation of treatment efficacy

After 2 and 6 w of treating CML, efficacy was observed in hematology, molecular biology and cytogenetics assays.

Hematology: Complete remission was defined as leukocyte $<10 \times 10^9/L$, platelet $<450 \times 10^9/L$ and less than 0.05 of primordial cells in bone marrow.

Cytogenetics: Bone marrow samples were collected to analyse 20 cells at metaphase by R band visualization. Complete remission was defined with absence of Ph chromosome positive cells at metaphase.

Molecular biology: PCR was used to measure *BCR/ABL* mRNA level. Complete remission was identified when PCR results were negative based on international standard of *BCR-ABL* gene transcription level [15].

Statistical analysis

SPSS 16.0 software was used for statistical analysis of all data, of which measurement data were presented as mean \pm Standard Deviation (SD). One-way Analysis of Variance (ANOVA) was used for comparison among multiple groups, followed by SNK-q test for between-group comparison. Enumeration data were analysed by chi-square test. Spearman approach was used for correlation analysis between two factors. A statistical significance was defined when $p < 0.05$.

Results

General information of patients

By analyzing general information of CML patients and healthy volunteers in clinics, we found no significantly statistical difference of gender, age and Body Mass Index (BMI) between two groups, which were thus comparable ($p > 0.05$, Table 2).

Bcl-2 and *c-myc* mRNA expression at different stages

Real-time fluorescent quantitative PCR has higher sensitivity and specificity than routine PCR. Using healthy population as the control group, *Bcl-2* and *c-myc* mRNA expressions were quantified relatively. Results showed that with elongated treatment time, *Bcl-2* mRNA level was significantly elevated after 6 w of combined treatment ($p < 0.05$ compared with those at admission). *C-myc* level was significantly elevated at w 2 and w 6 of combined treatment group. Single use of hydroxyurea remarkably elevated expression level at 6 weeks after treatment ($p < 0.05$ compared with those in admission, Figure 1). We further performed agarose gel electrophoresis on

Effects of recombinant human interferon $\alpha 1b$ on *Bcl-2* and *c-myc* expression in chronic myeloid leukemia and correlation with clinical treatment efficacy

amplified products. Images were shown in Figures 2 and 3, with consistent results as those from quantification assay.




Figure 1. *Bcl-2* and *c-myc* mRNA expression level at different stages during treatment. ** $p < 0.05$ comparing between combined treatment and healthy control; ## $p < 0.05$ comparing between hydroxyurea and healthy control group, && $p < 0.05$ comparing combined treated group and those at onset.




Figure 2. Electrophoresis of *c-myc* amplification bands.



Figure 3. Electrophoresis of *Bcl-2* amplification product.

Comparison of efficacy among different treatment methods

After 2 or 6 w of combined treatment or hydroxyurea treatment, efficacy was analysed from hematology, cytology and genetics. Results showed that combined treatment of recombinant human interferon $\alpha 1b$ plus hydroxyurea had better efficacy than singly treatment using hydroxyurea from all perspectives, as shown in Table 3.

Correlation between *Bcl-2/c-myc* mRNA expression level and treatment efficacy

Pearson analysis was performed to reveal the correlation between *Bcl-2* or *c-myc* mRNA expression and clinical treatment efficacy. Results showed positive correlation of treatment efficacy with *Bcl-2* mRNA expression ($r=0.541\sim 0.623$, $p < 0.01$), and *c-myc* mRNA expression ($r=0.645\sim 0.612$, $p < 0.01$, Table 4).

Table 1. *Bcl-2* and *c-myc* primer sequences.

Gene	Primer sequence	Length
β -actin	Forward: 5'-AAACTGGAACGGTGAAGGTG-3'	119 pb

Reverse: 5'-AGTGGGGTGGCTTTTAGGAT-3'

Bcl-2 Forward: 5'-CTCCCGCCGCCGCTACCGC-3' 125 pb

Reverse: 5'-CTGGGGCCGTACAGTTAC-3'

c-myc Forward: 5'-TTCTCTCCGTCCTCGGATTC-3' 116 pb

Reverse: 5'-GTAGTTGTGCTGATGTGTGGA-3'

Table 2. Generation information of participants.

Related index	CML	Control	P value
N	95	38	-
Gender			
Male (N)	57	22	0.832
Female (N)	38	16	
Age (years)	55.2 \pm 13.2	59.6 \pm 11.7	0.075
BMI (kg/m ²)	20.5 \pm 2.5	20.4 \pm 2.1	0.828

Table 3. Comparison of treatment efficacy among different methods.

Group	CHR	CMR	CCyR
Combined treatment	86.32%**	39.55%**	57.90%**
Hydroxyurea	60.86%	20.14%	37.54%

Note: CHR: Complete Hematology Reaction; CMR: Complete Molecular Response; CCyR: Complete Cytogenetics Response. ** $p < 0.05$ compared to hydroxyurea treatment group.

Table 4. Correlation between *Bcl-2/c-myc* mRNA expression and treatment efficacy.

Treatment efficacy	<i>Bcl-2</i>		<i>c-myc</i>	
	r	P	r	P
Hematology	-0.541	$P < 0.01$	-0.621	$P < 0.01$
Molecular biology	-0.562	$P < 0.01$	-0.645	$P < 0.01$
Cytogenetics	-0.623	$P < 0.01$	-0.612	$P < 0.01$

Discussion

Hydroxyurea exerts its treatment effects *via* inhibiting nucleic acid reductase to interfere with DNA synthesis on S phase [16,17]. It usually functions on late stage of hematological system, thus having minor suppression on bone marrow. Interferon $\alpha 1b$ has property of anti-virus and suppression of cell proliferation without selectivity. Therefore, it is more effective to use both hydroxyurea and interferon, but without clearly illustrated functional mechanisms.

Results of this study showed lower expression of tumor suppressor gene *Bcl-2* and *c-myc* in CML patients compared with control population, consistent with previous studies [18,19]. After using both hydroxyurea and interferon for treating CML patients, mRNA expression of *Bcl2* and *c-myc* was elevated with extended treatment time, indicating that

hydroxyurea and recombinant $\alpha 1b$ could exert treatment effects *via* facilitating expression of tumor suppressor gene. However, compared with single use of hydroxyurea, combined treatment can effectively accelerate *Bcl-2* and *c-myc* up-regulation. *In vitro* study of interferon α -bone marrow cell co-culture showed that interferon α could affect cell biological behavior *via* modifying *Bcl-2* or *c-myc* gene expression, thus alleviating disease severity. The results of *in vivo* study were consistent with those *in vitro* studies. Interferon α suppressed *BCR/ABL* gene expression in CML leukemia cell lineage [20]. *BCR/ABL* fusion gene can exert anti-apoptotic effects *via* enhancing expression of anti-apoptotic gene *Bcl-2*.

Combined treatment using both recombinant human interferon $\alpha 1b$ and hydroxyurea had better efficacy than hydroxyurea alone from perspectives of cytology, hematology and genetics. Combining with previous study data, using both α interferon and imatinib for treating CML had better efficacy than single use of imatinib in both low risk and moderate risk groups, but not in those high risk patients [21]. Alpha-interferon had certain diagnostic values for hairy cell leukemia and CML [22]. However, analysis of different CML grades was not performed in the present study and requires further investigation in the future. Interestingly, our study showed positive correlation between *Bcl-2/c-myc* expression and clinical treatment efficacy as demonstrated by improved treatment efficacy along with elevated *Bcl-2* or *c-myc* expression. However, due to limited numbers of patients enrolled in the present study, large cohort clinical studies are required to confirm these findings in the future.

Conclusion

Combined treatment of interferon and hydroxyurea had better efficacy than single use of hydroxyurea from cytology, hematology and genetics perspectives. Both treatments elevated mRNA expressions of *Bcl-2* and *c-myc*. Therefore, recombinant human interferon $\alpha 1b$ has treatment effects on CML.

References

- Jabbour E, Kantarjian H. Chronic myeloid leukemia: 2016 update on diagnosis, therapy, and monitoring. *Am J Hematol* 2016; 91: 252-265.
- Heisterkamp N, Stam K, Groffen J, de Klein A, Grosveld G. Structural organization of the *bcr* gene and its role in the Ph translocation. *Nature* 1985; 315: 758-761.
- Weissman IL. Stem cells: units of development, units of regeneration, and units in evolution. *Cell* 2000; 100: 157-168.
- Vinhas R, Cordeiro M, Pedrosa P, Fernandes AR, Baptista PV. Current trends in molecular diagnostics of chronic myeloid leukemia. *Leuk Lymphoma* 2016; 1-14.
- Kantarjian H, O'Brien S, Jabbour E, Garcia-Manero G, Quintas-Cardama A, Shan J, Rios MB, Ravandi F, Faderl S, Kadia T, Borthakur G, Huang X, Champlin R, Talpaz M, Cortes J. Improved survival in chronic myeloid leukemia since the introduction of imatinib therapy: a single-institution historical experience. *Blood* 2012; 119: 1981-1987.
- Jabbour E. Chronic myeloid leukemia: first-line drug of choice. *Am J Hematol* 2016; 91: 59-66.
- Baccarani M, Castagnetti F, Gugliotta G, Palandri F, Rosti G. Treatment recommendations for chronic myeloid leukemia. *Mediterr J Hematol Infect Dis* 2014; 6: e2014005.
- Talpaz M, Hehlmann R, Quintas-Cardama A, Mercier J, Cortes J. Re-emergence of interferon-alpha in the treatment of chronic myeloid leukemia. *Leukemia* 2013; 27: 803-812.
- Takeuchi K. Analysis of IFN-alpha inducible genes in human chronic myelogenous leukemia cell line KT-1. *Rinsho Ketsueki* 2003; 44: 65-69.
- Yassin MA, Kohla S, Al-Sabbagh A, Soliman AT, Yousif A, Moustafa A, Battah AA, Nashwan A, Al-Dewik N. A case of chronic neutrophilic leukemia successfully treated with pegylated interferon alpha-2a. *Clin Med Insights Case Rep* 2015; 8: 33-36.
- Willimott S, Wagner SD. Post-transcriptional and post-translational regulation of Bcl2. *Biochem Soc Trans* 2010; 38: 1571-1575.
- Chanvorachote P, Nimmannit U, Stehlik C, Wang L, Jiang BH, Ongpipatanakul B, Rojanasakul Y. Nitric oxide regulates cell sensitivity to cisplatin-induced apoptosis through S-nitrosylation and inhibition of Bcl-2 ubiquitination. *Cancer Res* 2006; 66: 6353-6360.
- Lin CY, Loven J, Rahl PB, Paranal RM, Burge CB, Bradner JE, Lee TI, Young RA. Transcriptional amplification in tumor cells with elevated *c-Myc*. *Cell* 2012; 151: 56-67.
- Milner AE, Grand RJ, Vaughan AT, Armitage RJ, Gregory CD. Differential effects of BCL-2 on survival and proliferation of human B-lymphoma cells following gamma-irradiation. *Oncogene* 1997; 15: 1815-1822.
- Hughes T, Deininger M, Hochhaus A, Branford S, Radich J, Kaeda J, Baccarani M, Cortes J, Cross NC, Druker BJ, Gabert J, Grimwade D, Hehlmann R, Kamel-Reid S, Lipton JH, Longtine J, Martinelli G, Saglio G, Soverini S, Stock W, Goldman JM. Monitoring CML patients responding to treatment with tyrosine kinase inhibitors: review and recommendations for harmonizing current methodology for detecting BCR-ABL transcripts and kinase domain mutations and for expressing results. *Blood* 2006; 108: 28-37.
- Kennedy BJ, Yarbro JW. Metabolic and therapeutic effects of hydroxyurea in chronic myeloid leukemia. *JAMA* 1966; 195: 1038-1043.
- de Lima PD, Cardoso PC, Khayat AS, Bahia Mde O, Burbano RR. Evaluation of the mutagenic activity of hydroxyurea on the G1-S-G2 phases of the cell cycle: an *in vitro* study. *Genet Mol Res* 2003; 2: 328-333.
- Cramer K, Nieborowska-Skorska M, Koptyra M, Slupianek A, Penserga ET, Eaves CJ, Aulitzky W, Skorski T. BCR/ABL and other kinases from chronic myeloproliferative disorders stimulate single-strand

Effects of recombinant human interferon $\alpha 1b$ on Bcl-2 and c-myc expression in chronic myeloid leukemia and correlation with clinical treatment efficacy

- annealing, an unfaithful DNA double-strand break repair. *Cancer Res* 2008; 68: 6884-6888.
19. Delgado MD, Leon J. Myc roles in hematopoiesis and leukemia. *Genes Cancer* 2010; 1: 605-616.
20. Yanagisawa K, Yamauchi H, Kaneko M, Kohno H, Hasegawa H, Fujita S. Suppression of cell proliferation and the expression of a BCR-ABL fusion gene and apoptotic cell death in a new human chronic myelogenous leukemia cell line, KT-1, by interferon-alpha. *Blood* 1998; 91: 641-648.
21. Bohn JP, Gastl G, Steurer M. Long-term treatment of hairy cell leukemia with interferon- α : still a viable therapeutic option. *Memo* 2016; 9: 63-65.
22. Burchert A, Saussele S, Eigendorff E, Muller MC, Sohlbach K, Inselmann S, Schutz C, Metzelder SK, Ziermann J, Kostrewa P, Hoffmann J, Hehlmann R, Neubauer A, Hochhaus A. Interferon alpha 2 maintenance therapy may enable high rates of treatment discontinuation in chronic myeloid leukemia. *Leukemia* 2015; 29: 1331-1335.

***Correspondence to**

Pengxiang Guo

Department of Hematology

Guizhou Provincial People's Hospital

PR China