

Environmental cues to urgency and incontinence episodes in Chinese patients with overactive urinary bladder syndrome.

Hongxia Su¹, Sen Yang¹, Yijun Wang¹, Min Chen², Aishu Dong³, Yuru Huang⁴, Cong Zhu²

¹Department of Urinary Surgery, ²Department of Nursing, ³Department of Cardiovascular, ⁴ICU, The Second Affiliated Hospital & Yuying Children's Hospital of Wenzhou Medical University, Wenzhou, Zhejiang, 325027, China.

Abstract

The present study was designed to investigate the possible environmental cues associated with urinary urgency and urinary incontinence in Chinese patients with diagnosed overactive bladder syndrome (OAB). A semistructured questionnaire was conducted with a convenience sample of 76 patients with OAB. Patients were interviewed about environmental cues associated with urgency and incontinence. The frequency of OAB episodes to each of the cues was quantified. Interviews lasted for seven to twenty-six minutes. Our results showed that 100% (n=76) respondents reported at least one cue associated with urgency and 92.1% (n=70) reported at least one cue associated with incontinence. "On the way to the washroom" frequency of the respondents was reported to be, 89.5% for urgency and 81.6% for incontinence. Our study, therefore suggested that one or more environmental cues might influence episodes of urinary urgency and incontinence in patients with OAB. A combination of behavioral treatments targeting environmental cues and Pavlovian extinction treatments may be effective in treating OAB.

Keywords: Environmental cues, urinary urgency, urinary incontinence, overactive bladder syndrome

Accepted February 09 2015

Introduction

Overactive bladder syndrome (OAB) is known to be a highly widespread disorder that increases with age in both sexes has negative effect on quality of life [1]. It was reported that approximately 6% of Chinese population suffer from OAB and other forms of incontinence [2].

According to the International Continence Society (ICS), OAB is a common disorder of bladder dysfunction, which is defined as urgency with or without urge incontinence, is usually with frequency and nocturia [3]. OAB is usually treated with nonpharmacologic methods such as lifestyle interventions (fluid restriction, avoidance of caffeine), bladder retraining and/or the pharmaceutical agents such as antimuscarinic drugs (e.g., darifenacin, hyoscyamine, oxybutynin) [4,5].

However, these treatments have proven effective only for some OAB symptoms. Behavioral treatments may not adequately relieve all bothersome symptoms [6]. In addition, pharmacological treatments are costly and have adverse side effects [7].

To date, the precise etiology of OAB remains poorly understood. OAB was reported to be associated with over

activity of the urinary bladder detrusor muscle and increased contraction that originates from the urothelium and lamina propria [8]. Recently, environmental cues are reported to be related to OAB symptoms [9,10]. The cue that is most mentioned was upon arrival at an individual's front door is defined as "latchkey incontinence", which is a loss of urine that occurs when one arrives home and puts the key in the lock of one's front door [9].

Study by Ghei and MaloneLee [11].showed that 52% sample of nearly 1800 patients with OAB had latchkey urgency and 26% had latchkey incontinence, suggesting that environmental cues, which as conditioned stimuli, play an important role in the development of OAB. Therefore, to identify possible environmental cues associated with OAB for treatment of urinary urgency and incontinence, environmental stimulus control training could be one of the useful methods.

Environmental cues to urgency and incontinence episodes in Chinese patients with overactive bladder syndrome have not been reported. Therefore, in the current study, a pilot study is conducted by planning structured interview techniques to investigate the possible environmental cues associated with urinary urgency and urinary incontinence in Chinese patients with diagnosed OAB.

Methods

Study Samples

Seventy-six (76) OAB patients recruited in the pilot study were diagnosed and treated in the Second Affiliated Hospital of Wenzhou Medical University between January, 2013 and March, 2014. Inclusion Criteria included, to literate patients with OAB based on the subject's score on the OAB Scale (OABSS) [12]. Participants were excluded if they had cognitive or other impairments that would interfere with completing the structured interview or refused to perform the study. This study was approved by the Institutional Review Board of The Second Affiliated Hospital of Wenzhou Medical University. Written informed consent was obtained from all participants.

Study procedures

All participants were interviewed at the urologist's office before or after the patient's scheduled appointment. Interview was conducted using semistructured questionnaire, including open- and closed-ended questions about environ-mental cues associated with urgency and incontinence. Patients were asked to respond to questions using a likert response rating scale: never, rarely, occasionally, often and always; which was coded from 1-5 for analysis. The frequency of OAB episodes to each of the cues was quantified. Interviews lasted between 7 and 26 minutes (mean 15 minutes).

Data Statistics

Descriptive statistics were computed for each cue for both urgency and incontinence situations. Quantitative data were

shown as median, qualitative data were shown as percentage or absolute value.

Results

Characteristics of Participants

Seventy-six (76) patients with OAB were recruited. Majority of participants (84.2%, n=64) were female. The median age was 36 years (range, 20 to 68 years) and most of the patients were married (80.3%, n=61). The educational level in our study samples was relatively low. 44.7% (n=34) completed

Table 1. The demographic characteristics of participants (n=76).

Characteristic	n (%) or median (range)
Gender	
Male	12 (15.8)
Female	64 (84.2)
Age (years)	36 (20-68)
OAB duration	3.5 (0.25-9)
Educational level	
Primary school (grades 1–6)	34 (44.7)
High school (grades 7–12)	26 (34.2)
University	16 (21.1)
Marital status	
Married	61 (80.3)
Single	11 (14.5)
Divorced or widowed	4 (5.3)

Table 2. Cues for urgency and incontinence (N = 76).

	Cues for urgency			Cues for incontinence		
	N	%	Frequency (1-5)	N	%	Frequency (1-5)
On the way to the washroom	68	89.5	4	62	81.6	3.5
Getting up after waking in the morning	65	85.5	4	56	73.7	3.5
Arrival home and opening the front door	54	71.1	4	52	68.4	3
Being cold	50	65.8	2.5	33	43.4	3
Arising from a lying-down position	44	57.9	3.5	27	35.5	3
Seeing running water	40	52.6	3.5	27	35.5	3
Think about going to the bathroom	28	36.8	4	10	13.2	4
In the bathroom for another reason besides urinating	20	26.3	3	7	9.2	5
After having tea	15	19.7	3	5	6.6	3
Knowing I will not be near a bathroom for a while	13	17.1	3	4	5.3	3
Removing my undergarments to take a shower/change clothes	12	15.8	3	4	5.3	3
Near a bathroom I often use	12	15.8	3	3	3.9	3
After having alcohol	10	13.2	3	1	1.3	2
Someone mentions going to the bathroom	8	10.5	3	0	0	0
Near a bathroom I occasionally use	6	7.9	2.5	2	2.6	3
After completing a task	5	6.6	3	1	1.3	3
Arising from a sitting position	3	3.9	3	1	1.3	3
Seeing signs/advertisements for bathrooms	3	3.9	2	1	1.3	2
Answering this questionnaire	2	2.6	2	0	0	0

primary school level, 34.2% (n=26) completed high school and only 21.1% (n=16) had bachelor or higher degree. The median disease duration was 3.5 years (range, 3 months to 9 years). Details are shown in Table 1.

Cues Survey Results

Table 2 shows the responses according to frequency of endorsement for urgency and for incontinence, respectively. 100% (n=76) reported at least 1 cue associated with urgency and 92.1% (n=70) reported at least 1 cue associated with incontinence.

The following cues were endorsed as leading to urgency by at least 50% of the participants. 1) On the way to the washroom, 2) getting up after waking in the morning, 3) arrival home and opening the front door, 4) being cold, 5) arising from a lying-down position, and 6) seeing running water. With the exception of "arising from a lying-down position, and seeing running water", the same cues were endorsed by at least 50% of the participants as associated with incontinence (Table 2).

The following cues were endorsed as leading to urgency with the median of frequency ≥ 3.5 . 1) On the way to the washroom, 2) getting up after waking in the morning, 3) arrival home and opening the front door, 4) arising from a lying-down position, 5) seeing running water, and 6) think about going to the bathroom. With the exception of "arrival home and opening the front door", "arising from a lying-down position", "seeing running water" was also evident. In addition, "In the bathroom for another reason besides urinating", the same cues were endorsed as associated with incontinence with the median of frequency ≥ 3.5 (Table 2).

Discussion

Results of our study indicate that patients diagnosed with OAB were associated with one or more environmental cues with urgency and incontinence. The most frequently reported were environmental cues associated with being on the way to a bathroom, getting up after waking in the morning, arrival home and opening the front door, arising from a lying-down position, seeing running water, thinking about going to the bathroom and in the bathroom for another reason besides urinating. These environmental cues either occur in OAB patients (at least 50%), or induce the episodes of urinary urgency and incontinence with high frequency (≥ 3.5). Therefore, the clinicians and patients should take into account of these cues and avoid them if conditions allowed.

Our results demonstrated that the environmental cues for OAB symptoms are those that occur in either temporal or spatial proximity to usual voiding situations. These findings support the hypothesis that Pavlovian condi-

tioning may influence OAB symptoms. Pavlovian conditioning has been found to underlie an extensive range of responses, from drug tolerance to emotional reactions [13]. Pavlovian conditioning occurs in nearly all animal systems and is believed to help organisms to prepare for physiologically significant events such as eating or fleeing a threat [14]. For example, frequent concurrence of immediate urination with arriving home leads to a conditioned response of a strong urge to urinate and actual loss of urine upon arriving home. Our results also indicated a discrepancy between the urinary urgency and incontinence symptoms induced by environmental cues. Some cues (such as arrival home and opening the front door) were more associated with the symptom of urgency, while others (such as on the way to the washroom) may contribute to sphincter incontinence. However, these findings need to be investigated.

Animal study indicates that extinction of Pavlovian conditioning is possible [15]. Therefore, new methods on extinction of Pavlovian conditioning could be extended to the treatment of the episodes of OAB through stimulus control training. For example, OAB patients should consciously choose other activities when arriving at home, or unlinking of the arrival at home with voiding. Moreover, OAB patients should avoid these conditioned stimuli, such as thinking other things, or taking different routes to the bathroom. However, systematic, admitted and general protocols for extinction of bladder responses to cues have yet to be developed and this is our next research direction and goals.

Limitations

There were several limitations in our study. First, the study sample was relative small and the results might limit the generalizability of the observations to the OAB patient population at large. Secondly, responses to environmental cues were dependent on the respondent's memory, thus the respondents may not be aware of the cues that actually affected their urgency and incontinence episodes. Thirdly, the methods that we applied were self-design semistructured questionnaire. Although the interview included open and closed-ended questions, it did not include all the possible environmental cues that were associated with urgency and incontinence. Finally, the frequency of endorsement for urgency and incontinence by OAB patients could not be statistically analyzed to obtain statistical differences. The limitation of our study requires further investigation to collect data from a much larger and diverse number of samples. Future studies might also investigate recognition of environmental cues to urgency and incontinence, which might help people with OAB to recognize conditions of the environment that exacerbate their symptoms.

Conclusion

In conclusion, the findings of the current study demonstrate that one or more environmental cues may influence episodes of urinary urgency and incontinence in patients with OAB. Pavlovian conditioning plays a role in OAB symptoms. A combination of behavioral treatments targeting environmental cues and Pavlovian extinction treatments might be effective in treating OAB.

Conflict of interest

The authors declare no conflict of interest.

References

1. Temml C, Heidler S, Ponholzer A, et al. Prevalence of the overactive bladder syndrome by applying the International Continence Society definition. *Eur Urol.* 2005; 48: 622-627.
2. Wang Y, Xu K, Hu H, et al. Prevalence, risk factors, and impact on health related quality of life of overactive bladder in China. *Neurourol Urodyn.* 2011; 30: 1448-1455.
3. Neveus T, von Gontard A, Hoebeke P, et al. The standardization of terminology of lower urinary tract function in children and adolescents: report from the Standardisation Committee of the International Children's Continence Society. *J Urol.* 2006; 176: 314-324.
4. Sussman DO. Overactive bladder: treatment options in primary care medicine. *J Am Osteopath Assoc.* 2007; 107: 379-385.
5. Radchenko VH. [Overactive bladder syndrome: etiology, pathogenesis, treatment]. *Fiziol Zh.* 2013; 59: 124-128.
6. Wyman JF, Harding G, Klutke C, et al. Contributors to satisfaction with combined drug and behavioral therapy for overactive bladder in subjects dissatisfied with prior drug treatment. *J Wound Ostomy Continence Nurs.* 2010; 37: 199-205.
7. Arnold J, McLeod N, Thani-Gasalam R, et al. Overactive bladder syndrome - management and treatment options. *Aust Fam Physician.* 2012; 41: 878-883.
8. Sacco E. [Physiopathology of overactive bladder syndrome]. *Urologia.* 2012; 79: 24-35.
9. Victor E, O'Connell KA, Blaivas JG. Environmental cues to urgency and leakage episodes in patients with overactive bladder syndrome: a pilot study. *J Wound Ostomy Continence Nurs.* 2012; 39: 181-186.
10. O'Connell KA, Torstrick A, Victor E. Cues to urinary urgency and urge incontinence: how those diagnosed with overactive bladder syndrome differ from undiagnosed persons. *J Wound Ostomy Continence Nurs.* 2014; 41: 259-267.
11. Ghei M, Malone-Lee J. Using the circumstances of symptom experience to assess the severity of urgency in the overactive bladder. *J Urol.* 2005; 174: 972-976.
12. Blaivas JG, Panagopoulos G, Weiss JP, et al. Validation of the overactive bladder symptom score. *J Urol.* 2007; 178: 543-547.
13. Wagner AR. Evolution of an elemental theory of Pavlovian conditioning. *Learn Behav.* 2008; 36: 253-265.
14. Delamater AR. Issues in the extinction of specific stimulus-outcome associations in Pavlovian conditioning. *Behav Processes.* 2012; 90: 9-919.
15. Bouton ME. Context and behavioral processes in extinction. *Learn Mem.* 2004; 11: 485-494.

*Correspondence to:

Cong Zhu
Department of Nursing
The Second Affiliated Hospital of Wenzhou Medical University
109, West Xueyuan Road, Wenzhou
Zhejiang Province, 325007
China.