

Vestibular function and curative effect of BPPV patients in different duration by application of electronystagmography and posturography.

Jianguo Liu[#], Xubo Chen[#], Ziwei Hu, Meiqun Wang, Wen Xie, Suying Guo, Yuehui Liu^{*}

Department of Otorhinolaryngology, the Second Affiliated Hospital of Nanchang University, Jiangxi Province, China

[#]These authors contributed equally to this study

Abstract

Objects: Benign paroxysmal positional vertigo (BPPV) is a common peripheral vestibular disease clinically. Electronystagmography (ENG) and posturography are commonly used methods for diagnosis of BPPV and examination of the vestibular function. Sparse study reported the characteristics of nystagmus and vestibular function and the curative effect in different duration of BPPV patients.

Methods: This is a retrospective study of 40 adults with vestibular symptoms. Before ENG test and posturography test, the subjects underwent physical examination and excluded central disorders. The canalith repositioning maneuver was used for the therapy of BPPV and the curative effect was studied between different duration groups.

Results: Of patients who suffered BPPV, higher rates of abnormal ENG were observed in the longer duration of BPPV. Among patients of BPPV, the patients' vestibular function in more than 12 months duration was inferior to the patients of a shorter duration. Moreover, the curative effect by repositioning maneuver in the shorter duration was superior to the longer duration.

Conclusions: The present study suggested that the vestibular function in patients that suffered BPPV deteriorates as the course of BPPV develops. The curative effect by repositioning maneuver in the patients of shorter duration is more satisfied than the longer duration.

Keywords: Vestibular function tests, Vertigo, Benign paroxysmal positional vertigo, Electronystagmography, Posturography.

Accepted on January 14, 2019

Introduction

With regards to peripheral vertigo, benign paroxysmal positional vertigo (BPPV) accounting for the major causes [1]. BPPV occurs due to the detachment of calcium-carbonate crystals or otoconia within the fluid-filled semicircular canals of the inner ear [2,3]. The clinical symptoms of BPPV varied among patients, symptoms may manifest with nonspecific dizziness, postural instability, mild headache, nausea and vomit [4,5]. However, the common manifestation is nystagmus in multiple styles. Deficit in different semicircular canals characterized by specified nystagmus in specified posture and is helpful in guiding canalith repositioning therapy especially in refractory cases with atypical BPPV [6,7].

Idiopathic BPPV is commonly occurred in the posterior and horizontal semicircular canals [8] while rarely onset in the anterior semicircular canal. Meanwhile, BPPV arising from multiple canals has also been described previously [9], the most common is a combination of posterior and horizontal BPPV as a constitution of 1.5%-5% in all BPPV patients [9,10]. Trauma may be a potential risk to cause combined canal BPPV but not a risk for bilateral single-canal BPPV [10].

Additionally, BPPV is more likely to develop in the right ear, a factor was reported to involve with the habit of sleeping on the right side in the major public [11].

In clinic, each type of BPPV is frequently diagnosed by observing the type of nystagmus induced by different positioning maneuvers, which were also the commonly used methods for restoration. The three widely used maneuvers including Semont's maneuver, for the posterior canal BPPV; Epley's maneuver, for the anterior canal BPPV; and Brandt-Daroff exercises, for the lateral canal BPPV. Some disorders such as posterior circulation stroke can mimic BPPV. However, central positional nystagmus usually manifested with sustaining vertigo and disequilibrium and other neurological symptoms and signs [12]. Presently, electronystagmography (ENG) is a common approach to differentiate vertigo and evaluate the effect after repositioning. Nonetheless, a few literatures compared the characteristic of the oculomotor system between different duration of BPPV before and after rehabilitation.

The visual system and proprioception also play negligible roles in postural control. Approximately 80% of an individual's sensory perception is gathered by the visual system [13], which

processes and integrates other sensory inputs, such as from vestibular, to select a balancing strategy [14]. Vestibular system is another crucial part for sustaining balance. Vestibular oculomotor reflex and vestibulospinal reflex are two major nervous reflex activities involving maintaining certain posture. Therefore, it is significant for diagnosis of BPPV by examination of static and dynamic posturography, which is a quantitative assessment of postural sway during stance. Though physicians are customizing treatments for their patients based on the information from posturography, to our knowledge, few studies reported the traits and curative effects of BPPV patients in different durations.

In this study, we analysed the traits of electronystagmography in patients with BPPV, studied the characteristic in different duration of BPPV before and after rehabilitation. Moreover, we explored the peculiarity of parameters in the equilibrium system in BPPV patients by posturography analysis. Our present study will throw light on diagnosis and proper administration of BPPV in clinic.

Materials and Methods

This was a descriptive, cross-sectional and observational study performed without intervention. Literary consent was mandatory for each subject. The present study was conducted with approval granted by the Committee on Clinical Research of Jiangxi Medical College, Nanchang University. Patients in the present study were recruited in outpatient department and inpatient department of otorhinolaryngology and neurology and even orthopedic during March 2008 to April 2011 in our hospital. The recruited patients were diagnosed with idiopathic BPPV based on the medical history and Dix-Hallpike test as well as Roll test. Routine otoscopic examinations were conducted for each patient in this study to exclude patients suffered with otitis media. Moreover, the central disorders were excluded in each object. The control subjects were recruited from outpatient by voluntary agreement.

Visual electronystagmography test

Electronystagmogram instrument equipped with introducing heated and cooled air was used to detect the Temperature experiment, saccadic movement, smooth pursuit test, gaze test and optokinetic nystagmus test in the patients.

Posturography test

Static posturography (SPG) and dynamic posturography (DPG) were included. The SPG quantifies sway velocity in degrees in each second ($^{\circ}/s$) with the participant standing in four test conditions: a firm surface with the eyes open; a firm surface with the eyes closed; a foam surface with eyes open and a foam surface with eyes closed. Twenty second trials were recorded for each test conditions. The average sway velocity of the participant's center of gravity was calculated per trial and averaged for each test condition. Falling down, crabbing rail or making a step during the test were recognized as an invalid test and make a fresh start. By contrast, DPG involves the use of

external balance perturbations or changing surface and/or visual conditions. Both the SPG and the DPG under each test condition were observed in a continuous 20 seconds.

Statistical analysis

In order to test the difference between two categorical variables, the χ^2 test was applied. All tests used in this study were 2-tailed, and a P value less than 0.05 were considered statistically significant. All statistical analyses were performed with SPSS Statistical software (version 20, Chicago, Illinois, USA).

Results

Totally 40 patients diagnosed with idiopathic BPPV were recruited, including 6 males and 34 females. The age of the patients was 34 to 79 years old. The duration from attacking with BPPV to seeking a medical advice in these patients was 1 week to 10 years. The duration of BPPV course was defined as the lasting time from the first occurrence of vertigo to ENG test in the study. The objects were matched in age and sex. We evaluated the ENG by stimulation of saccadic movement, smooth pursuit test, optokinetic nystagmus test, gaze test, spontaneous nystagmus and temperature experiment among control group, below 12 months course group (<12 months) of BPPV and above 12 months course group (>12 months). According to our data (Table 1), the saccadic movement, the smooth pursuit test and the optokinetic nystagmus test as well as the gaze test were in a normal condition in the control group and <6 months group. In the group of below 6 months course, all the patients detected spontaneous nystagmus by ENG. In the opposite, in the group of above 12 months course all patients not detected the spontaneous nystagmus (Table 1). As shown in the temperature experiment (Table 1), the below 6 months course group demonstrated a 35% positive rate and showed statistically significant higher positive rate when compared with the control group ($p < 0.05$). A higher positive rate was detected in the above 12 months course group in contrast with the below 6 months course group ($P < 0.05$).

Most vertigo complaints in clinic belong to peripheral vertigo, usually refers to disorder in vestibule. Other than ENG, dynamic posturography (DPG) is another valuable approach to evaluate the vestibular function. In the present study, by using DPG, proprioceptive score, visual score, vestibular score as well as Romber's score and comprehensive evaluation were assessed. As demonstrated in our study (Table 2), among the control group, below 6 months course group and above 12 months course group, these scores in DPG not showed a significant difference except the vestibular score.

Visual ocular system (VOS) test is also commonly used to evaluate the vestibular function, we compared the vestibular condition among control group, below 4 month course group (<4 months) and above 4 month course group (>4 months) in BPPV patients. Our data show that the positive rate in VOS test and temperature experiment was 15% and 10%, respectively. In <4 weeks group, the positive rate was 20% and

Vestibular function and curative effect of BPPV patients in different duration by application of electronystagmography and posturography

30%, respectively, while in >4 weeks group, the positive rate was ascended to 35% and 85%, respectively (Table 3).

At present, canalith repositioning maneuvers is the major therapeutical approach of BPPV. In order to investigate the effect of canalith repositioning maneuvers for treating BPPV in different BPPV course, we compared the curative effect among patients of <4 weeks group and >4 weeks group. There are 20 cases were studied in each group. After the treatment of canalolith repositioning maneuver, all the patients were visited by telephone follow-up for 3 months. Vertigo symptoms in patients accepted canalith repositioning maneuver occurred with lower frequency onset compared with previous duration

before the canalith repositioning maneuver were defined as improved. Vertigo symptom not alleviated after the canalith repositioning maneuver was defined as non-effective. The effect was divided into completely cured, improved and non-effective according to the degree of recovery. As the data demonstrated, the completely cured rate in <4 weeks group was higher than in >4 weeks group (Table 4). The effective rate, total of completely cured rate and improved rate, was higher in the <4 weeks group compared with the >4 weeks group. The rate of non-effective in the >4 weeks group is higher than the <4 weeks group (Table 4).

Table 1. Rates of abnormal ENG in different duration of BPPV patients (%).

Group	ENG							
	Saccadic movement	Smooth test	pursuit	Optokinnetic test	nystagmus	Gaze test	Spontaneous nystagmus	Temperature experiment
Control	0	0		0		0	0	10%
<6 months	0	0		0		0	100%	35% [#]
>12 months	100%	100%		100%		0	0	65% ^{*,+}

[#]P<0.05 when compared with control, ^{*}P<0.05 when compared with control group, ⁺P<0.05 when compared with <6 months group. Control group, n=20; <6 months group, n=14; >12 months group, n=26.

Table 2. Scores determined by DPG in patients with different courses of BPPV at the AP situation.

Group	DPG				
	Proprioceptive score	Visual score	Vestibular score	Romber's score	Comprehensive score
Control	91.62 ± 1.35	84.37 ± 1.92 [*]	62.91 ± 2.83	54.13 ± 1.61	69.55 ± 3.95
<6 months	92.58 ± 1.16	86.19 ± 2.99	57.63 ± 2.16 [#]	54.96 ± 2.18	67.43 ± 2.87
>12 months	93.77 ± 3.51	89.07 ± 3.67	51.98 ± 1.94 ^{#+}	57.03 ± 2.87	63.02 ± 4.89

[#]P<0.05 when compared with the control group; ^{*}P<0.05 when compared with the control group; ⁺P<0.05 when compared with the below 6 months course group. Data was shown as mean ± sem. Control group, n=20; <6 months group, n=14; >12 months group, n=26.

Table 3. Rate of abnormal vestibular function in different course of BPPV.

Group	Vestibular function	
	Positive rate in VOS	Positive rate in temperature experiment (%)
Control	15	10
<4 weeks	20.00 ^a	30.00 ^d
>4 weeks	35.00 ^{b,c}	85.00 ^{e,f}

^aP<0.05 when compared with the control group; ^bP<0.05 when compared with the control group; ^cP<0.05 when compared with the control group; ^dP<0.05 when compared with the control group; ^eP<0.05 when compared with the control group; ^fP<0.05 when compared with the control group. Control group, n=20; <4 weeks group, n=20; >4 weeks group, n=20.

Table 4. Analysis of curative effect of BPPV in different course.

Curative effect	<4 weeks	>4 weeks
-----------------	----------	----------

	Case	Rate (%)	Case	Rate (%)
Completely cured	18	90%	9	45%
Improved	2	10%	7	35%
Non-effective	0	0%	4	20%
Effective	20	100%	16	80%

Discussion

In clinic, BPPV is not difficult to be diagnosed based on the patient's medical history and physical examination [8]. Vestibule in normal function is crucial in maintaining body balance. Besides, other systems such as reactive, anticipatory, sensory, dynamic, and limits of stability and physiological systems such as vestibular, visual, proprioceptive, muscle strength, and reaction time contribute to balance. Thus, in these years, some assessment tools are presented and used to

evaluate balance, such as the Tinetti balance and gait test [15], the Berg balance scale [16], ENG [17], DPG and SPG [18].

In our present study, we found that the control patient and the <6 months course patient of BPPV not appeared nystagmus by saccadic movement test, smooth pursuit test and optokinetic nystagmus test (Table 1). However, in the >12 months course patient of BPPV all appeared with nystagmus by these tests. In addition, as indicated by DPG (Table 2) and VOR test (Table 3), the vestibule function score was descended as the disease of BPPV prolonged. These data may suggest that the vestibular function is worse in the longer course patient of BPPV. This may be explained by that the visual-vestibular optokinetic reflex is somewhat in deficits as the disease develops. The visual-vestibular optokinetic reflex is the primary mechanism for visual stabilization during head movements. Dysfunction of this reflex could arouse retinal slip occurs with head movement greater than approximately 2 Hz or 80 per second and visual acuity declines [19]. Cold or hot water/gas can be used to test whether any vestibular function after horizontal canal stimulation remains in the ear. In the temperature experiment, more cases were observed nystagmus in the longer course of BPPV (Table 1). The vestibule was irrigated by endolymphatic flow and nystagmus was occurred. However, not all patients of BPPV appeared with nystagmus by this test and our study was similar with previous study [20].

The prognosis of BPPV is excellent. As show in our data (Table 4), all of the patients <4 weeks course is recovered or partially recovered after a single maneuver, while in the patients >4 weeks course, 80% patients were recovered or improved. Our data is compatible with previous study which reported that 80-90% of the patients do not show signs of BPPV after the first treatment of canalith repositioning maneuver [21]. Except for canalith repositioning maneuver, some other approach such as chemical or even operational method is also applied for BPPV therapy in clinic [8,22]. As reported by Olusesi and colleague, vestibular suppressant was 68.4% success, particle repositioning maneuver was 69% success, and chemical labyrinthectomy was 53.8% success [23]. Therefore, canalith repositioning maneuver is the most effective and common method currently. Our data suggested that the patients with >4 week course may need multiple maneuver time [24] or some other therapeutic method. As reported by Korkmaz and coworker, age, sex, canal type and the duration of symptoms had no impact on the number of maneuvers [25]. However, as reported by Yoon and colleagues, longer duration of vertigo before treatment, bilateral or multiple canal involved, and age >50 years were the risk factors associated with the need for multiple particle repositioning maneuver [26]. These discrepancies may aroused by race of the patients.

In conclusion, our study suggested that the vestibular function in patients that suffered BPPV in a long duration was inferior to the shorter one. The curative effect by canalith repositioning maneuver in the patients of shorter duration is more satisfied than in the patients of longer suffering of BPPV.

Acknowledgment

The authors thank subjects in this study from the second affiliated hospital of Nanchang University.

Conflicting Interests

The authors declared no potential conflicts of interest to this study, authorship, and/or publication of this article.

Funding

This work was supported by the following fundings: 1) Science and Technology Support Program by Health and Family Planning Commission of Jiangxi Province (No. 20155260); 2) Science and Technology Support Program by Education Office of Jiangxi Province (No. GJJ14082); 3) Youth science foundation by Jiangxi Provincial Department of Science and Technology (No. 20142BBG70075v).

References

1. Somefun OA, Giwa OS, Bamgboye BA, Okeke-Igbokwe II, Azeez AA. Vestibular disorders among adults in a tertiary hospital in Lagos, Nigeria. *Eur Arch Otorhinolaryngol* 2010; 267: 1515-1521.
2. Hall SF, Ruby RR, McClure JA. The mechanics of benign paroxysmal vertigo. *J Otolaryngol* 1979; 8: 151-158.
3. Hamann KF. Benign paroxysmal positioning vertigo: a disease explainable by inner ear mechanics. *ORL J Otorhinolaryngol Relat Spec* 2006; 68: 329-333.
4. Blatt PJ, Georgakakis GA, Herdman SJ, Clendaniel RA, Tusa RJ. The effect of the canalith repositioning maneuver on resolving postural instability in patients with benign paroxysmal positional vertigo. *Am J Otol* 2000; 21: 356-363.
5. Giacomini PG, Alessandrini M, Magrini A. Long-term postural abnormalities in benign paroxysmal positional vertigo. *ORL J Otorhinolaryngol Relat Spec* 2002; 64: 237-241.
6. Aw ST, Todd MJ, Aw GE, McGarvie LA, Halmagyi GM. Benign positional nystagmus: a study of its three-dimensional spatio-temporal characteristics. *Neurology* 2005; 64: 1897-1905.
7. Lee SH, Kim JS. Benign paroxysmal positional vertigo. *J Clin Neurol* 2010; 6: 51-63.
8. Parnes LS, Agrawal SK, Atlas J. Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ* 2003; 169: 681-693.
9. Moon SY, Kim JS, Kim BK, Kim JI, Lee H, Son SI, Kim KS, Rhee CK, Han GC, Lee WS. Clinical characteristics of benign paroxysmal positional vertigo in Korea: a multicenter study. *J Korean Med Sci* 2006; 21: 539-543.
10. Tomaz A, Gananca MM, Gananca CF, Gananca FF, Caovilla HH, Harker L. Benign paroxysmal positional vertigo: concomitant involvement of different semicircular canals. *Ann Otol Rhinol Laryngol* 2009; 118: 113-117.

11. von Brevern M, Seelig T, Neuhauser H, Lempert T. Benign paroxysmal positional vertigo predominantly affects the right labyrinth. *J Neurol Neurosurg Psychiatry* 2004; 75: 1487-1488.
12. Kattah JC, Talkad AV, Wang DZ, Hsieh YH, Newman-Toker DE. HINTS to diagnose stroke in the acute vestibular syndrome: three-step bedside oculomotor examination more sensitive than early MRI diffusion-weighted imaging. *Stroke* 2009; 40: 3504-3510.
13. Friedrich M, Grein HJ, Wicher C, Schuetze J, Mueller A, Lauenroth A. Influence of pathologic and simulated visual dysfunctions on the postural system. *Exp Brain Res* 2008; 186: 305-314.
14. Schmid M, Nardone A, De Nunzio AM. Equilibrium during static and dynamic tasks in blind subjects: no evidence of cross-modal plasticity. *Brain* 2007.
15. Tinetti ME. Performance-oriented assessment of mobility problems in elderly patients. *J Am Geriatr Soc* 1986; 34: 119-126.
16. Berg K, Wood-Dauphinee S, Williams JI. The Balance Scale: reliability assessment with elderly residents and patients with an acute stroke. *Scand J Rehabil Med* 1995; 27: 27-36.
17. Gofrit SG, Mayler Y, Eliashar R, Bdolah-Abram T, Ilan O, Gross M. The association between vestibular physical examination, vertigo questionnaires, and the electronystagmography in patients with vestibular symptoms. *Ann Otol Rhinol Laryngol* 2017; 126: 315-321.
18. Mancini M, Horak FB. The relevance of clinical balance assessment tools to differentiate balance deficits. *Eur J Phys Rehabil Med* 2010; 46: 239-248.
19. Grossman GE, Leigh RJ, Bruce EN, Huebner WP, Lanska DJ. Performance of the human vestibuloocular reflex during locomotion. *J Neurophysiol.* 1989; 62: 264-272.
20. Batuecas-Caletrio A, Montes-Jovellar L, Boleas-Aguirre MS, Perez-Fernandez N. The ice-water caloric test. *Acta Otolaryngol* 2009; 129: 1414-1419.
21. von Brevern M, Seelig T, Radtke A, Tiel-Wilck K, Neuhauser H, Lempert T. Short-term efficacy of Epleys manoeuvre: a double-blind randomised trial. *J Neurol Neurosurg Psychiatry* 2006; 77: 980-982.
22. Volkenstein S, Dazert S. Recent surgical options for vestibular vertigo. *GMS Curr Top Otorhinolaryngol Head Neck Surg* 2017; 16.
23. Olusesi AD, Abubakar J. 10 years of vertigo Clinic at National Hospital Abuja, Nigeria: what have we learned? *Eur Arch Otorhinolaryngol* 2016; 273: 3567-3572.
24. Kansu L, Avci S, Yilmaz I, Ozluoglu LN. Long-term follow-up of patients with posterior canal benign paroxysmal positional vertigo. *Acta Otolaryngol* 2010; 130: 1009-1012.
25. Korkmaz M, Korkmaz H. Cases requiring increased number of repositioning maneuvers in benign paroxysmal positional vertigo. *Braz J Otorhinolaryngol* 2016; 82: 452-457.
26. Yoon J, Lee JB, Lee HY, Lee BD, Lee CK, Choi SJ. Potential risk factors affecting repeated canalith repositioning procedures in benign paroxysmal positional vertigo. *Otol Neurotol* 2018; 39: 206-211.

***Correspondence to**

Yuehui Liu
Department of Otorhinolaryngology
The second affiliated hospital of Nanchang University
Jiangxi Province
China